




INTEGRATED SYSTEM FOR PROTECTION OF BIRDS

Summary of Activities and the Results of Ornithological Monitoring in the Integrated System for Protection of Birds, 2021



Dr Pavel Zehtindjiev
Institute of Biodiversity and Ecosystem
Research-Bulgarian Academy of Sciences

Dr D. Philip Whitfield
Natural Research Ltd, Banchory, UK

Contents

Introduction	3
Results	5
Monitoring of geese in Winter 2020-2021	5
Spring migration	9
Autumn migration.....	13
List of participants in the observations	18

Introduction

Integrated System for Protection of Birds (ISPB) includes 114 wind turbines, 95 of which are within the Kaliakra SPA BG0002051 and 19 are in the areas adjacent to the protected zone.

The ornithological monitoring of ISPB is a complex study assigned by the Wind farms, located in Kaliakra SPA: BG0002051-AES Geo Energy Ltd., Kaliakra Wind Power, Degrets OOD, Disib OOD, Windex OOD, Long Man Invest OOD, Long Man Energy OOD, Zevs Bonus OOD, Vertikal-Petkov & Sie SD, Wind Park Kavarna East EOOD, Wind Park Kavarna West EOOD, Millennium Group OOD in 2021.

The ISPB consists of a combination of radar observations and meteorological data, integrated with field visual observations, which jointly used are essential for the accurate risk assessment and ensure that appropriate action is taken immediately to avoid collision risk. So far as potential adverse impacts of turbine collisions on birds, a Turbine Shutdown System is deployed supported by an Early Warning System.

The monitoring studies are based on the requirements of basic normative and methodological documents as follows: Environmental Protection Act, Biological Diversity Act, Bulgarian Red Data Book, Directive 92/43/EEC for habitats and species, and Directive 2009/147/EC on the conservation of wild birds, Protected Areas Act and Order RD-94 of 15.02.2018 of the Minister of Environment and Waters. Best international practices are also incorporated (T-PVS/Inf (2013) 15: <https://rm.coe.int/1680746245>). Detailed information on the scope, technical rules and monitoring procedures are publicly available at a dedicated website <https://kaliakrabirdmonitoring.eu/>.




Figure 1. A satellite photo with the location of the wind turbines covered by the ISPB and the boundaries of Kaliakra SPA (shown by the red line), together with the scope of three radar systems.

In order to provide objective data for the bird risk assessment, this summary presents activities and results of the monitoring in 2021.

The activities were supervised and coordinated by Prof. Dr. Pavel Zehtindjiev - Ornithologist with over 25 years of research in ornithology; over 85 scientific publications in international ornithological journals; member of European Ornithologists Union and several other conservation organisations; winner of the Revolutionary Discovery Award for Ornithology of an American Ornithological Society in 2016 – The Cooper Ornithological Society; more than 10 years of experience in impact monitoring of wind turbines on breeding, migrating and wintering bird species in the region of Kaliakra.

Three types of radars integrated into the ISPB were used for monitoring and prevention of bird collisions:

Bird Scan MS1

The radar collects quantitative data and provides information about Migration Traffic Rate of birds through a specific sector where the fixed beam of the radar is directed (Figure 1). The quality of the data deepens on the distance to the birds and to the size of the migrating birds. In the case of ISPB the maximum distance we have used the Bird Scan MS1 radar is 10 km beam directed from west to east across the main migratory front of seasonal migrations. The data obtained by this radar system allow crude identification of ecological types of birds: for example, passerines, swifts, waders and large birds. The radar data do not allow quantification of bird migration for every bird species observed in the ISPB territory and therefore do not allow any comparison with visual observations.

These data are not used for quantification and analysis of the characteristics of migration.

Deltatrack Radar System

This radar is a tracking radar system which allows detection of a single target or group of targets and tracking of their movements in a range of around 5 km (Figure 1). It is used in the monitoring as a real time tool for the tracking of already (visually) identified bird targets in the ISPB territory. The radar is not applicable for quantitative analysis of bird migration.

Radar System Robin

This is a 3-D radar system constructed for detection and tracking of moving targets in an air volume of around 10 km³ (<https://www.youtube.com/watch?v=-Kb70clGHOQ&t=8s>) (Figure 1). It is a real time tool for tracking of moving targets and in combination with visual observations in the field provides highly reliable data on the distance as well altitudes of birds already detected and identified by the field ornithologists. This radar does not provide quantitative data of migration at a species level because it does not allow species identification.

All three radar systems have been used as tools to assist field observations, detection of potential ingresses, and real time tracking of birds after visual observation through the ISPB during the period of monitoring.

All quantitative data and analysis of recorded bird numbers are based on the only possible quantification of bird migration of different bird species – the visual observations in the field. Locations of field observation points are presented in Figure 1 (white dots).

Detailed descriptions of the technical characteristics of the three radar systems integrated within the ISPB are presented on the web site: <http://kaliakrabirdmonitoring.eu/Methodology> .

Results

Monitoring of geese in Winter 2020-2021

The 90 days of the study encompassed the whole period when geese were recorded in the region during 2020-2021.

Total number of observed goose species and their numbers

In total very low numbers of geese of all observed species were present in the ISPB territory during the winter 2020-2021. Unusually low numbers of wintering geese were also observed in Bulgaria and Romania in general in the winter season 2020-2021 (<https://greenbalkans.org/bg/>, <https://BirdLife Bulgaria red brested geese>).

Less than 2000 individual geese were observed during the surveys (Table 1).

Table 1. The number of observed geese by dates of different species (data from visual observations). The dates with zero observed birds are not included. Species involve Greater White-fronted Goose (GWFG: *Anser albifrons*), Greylag Goose (*Anser anser*) and Red-breasted Goose (*Branta ruficollis*).

Date	A. albifrons	A. anser	Anser/Branta	Total
21.12.2020	110			110
02.01.2021	80			80
28.01.2021	668	24	975	1667
13.02.2021	42			42
Grand Total	900	24	975	1899

The four days with observed geese in ISPB territory are presented in Table 1. The maximum number of geese in flocks was observed on 28 January. There was only one day with geese observed in February. No Red-breasted Goose (RBG) was observed in February.

Temporal dynamics of geese number during the period when geese were observed in ISPB territory are presented in Figure 2.




Figure 2. Temporal dynamics of wintering geese observed in ISPB territory, season 2020-2021.

Spatial distribution of feeding geese in the ISPB territory

The flocks of geese tracked and confirmed visually are presented in maps below. Due to the low number of wintering geese in this winter spatial analysis was not possible. In general, (the same as in previous winters) concentration of flights in NE part of the ISPB territory was observed in winter 2020-2021. More detailed analysis of the feeding preferences of wintering geese in ISPB territory are presented in previous reports available at the web site of ISPB

https://kaliakrabirdmonitoring.eu/Report_Winter_2018-2019,

https://kaliakrabirdmonitoring.eu/Report_Winter_2019-2020,

https://kaliakrabirdmonitoring.eu/Report_Winter_2020-2021.




Figure 3. Flocks of of GWFG (yellow), Greylag goose (green) and mixed flocks of GWFG and RBG (blue) (yellow) observed during the monitoring period in winter 2020-2021 in ISPB territory.

Carcass monitoring results

All 114 turbines were programmed to be searched every seventh day in the periods of autumn and spring migration as well as during the wintering period of geese. The rest of the time during the whole year every turbine was searched once per month if the areas under turbines were accessible. During the winter monitoring (subject of this report) all 114 turbines were searched for carcasses during the whole winter survey period (01 December 2020 –28 February 2021) when more birds were at risk of collision. The frequencies of searches are presented in Table 2.

Table 2. Number of searches per turbine during the winter monitoring 2020-2021

Turbine code	December	January	February	Total
ABBalgarevo	2	4	4	10
ABГ1	2	4	4	10
ABГ2	2	4	4	10

Turbine code	December	January	February	Total
ABГ3	2	4	4	10
ABГ4	2	4	4	10
ABMillenium group	2	4	4	10

Turbine code	December	January	February	Total
ABMillenium group				
Mikon	2	4	4	10
AE10	2	4	4	10
AE11	2	4	4	10
AE12	2	5	4	11
AE13	2	5	4	11
AE14	2	3	4	9
AE15	2	5	4	11
AE16	2	4	4	10
AE17	2	4	4	10
AE18	2	5	4	11
AE19	2	5	4	11
AE20	2	4	4	10
AE21	2	4	4	10
AE22	2	4	4	10
AE23	2	4	4	10
AE24	2	4	4	10
AE25	2	4	4	10
AE26	2	4	4	10
AE27	2	4	4	10
AE28	2	4	4	10
AE29	2	4	4	10
AE31	2	5	4	11
AE32	2	5	4	11
AE33	2	5	4	11
AE34	2	5	4	11
AE35	2	5	4	11
AE36	2	4	4	10
AE37	2	5	4	11
AE38	2	4	4	10
AE39	2	4	4	10
AE40	2	4	4	10
AE41	2	4	4	10
AE42	2	4	4	10
AE43	2	4	4	10
AE44	2	4	4	10
AE45	2	4	4	10
AE46	2	5	4	11
AE47	2	5	4	11
AE48	2	5	4	11
AE49	2	5	4	11
AE50	2	5	4	11
AE51	3	4	4	11
AE52	3	4	4	11

Turbine code	December	January	February	Total
AE53	3	4	4	11
AE54	3	4	4	11
AE55	3	4	4	11
AE56	3	4	4	11
AE57	3	4	4	11
AE58	3	4	4	11
AE59	3	4	4	11
AE60	2	5	4	11
AE8	2	4	4	10
AE9	2	4	4	10
DBG1	2	4	4	10
DBG1HSW250	2	4	4	10
DBG2	2	4	4	10
DBG2MN600	2	4	4	10
DBG3	2	4	4	10
DBG4	2	4	4	10
DBG5	2	4	4	10
DC1	2	4	4	10
DC2	2	4	4	10
E00	2	4	4	10
E01	2	4	4	10
E02	2	4	4	10
E04	2	4	4	10
E05	2	4	4	10
E07	2	4	4	10
E08	2	4	4	10
E09	2	4	4	10
M1	2	4	4	10
M10	2	4	4	10
M11	2	4	4	10
M12	2	5	4	11
M13	2	5	4	11
M14	2	5	4	11
M15	2	5	4	11
M16	2	5	4	11
M17	2	5	4	11
M18	2	5	4	11
M19	2	5	4	11
M2	2	4	4	10
M20	3	4	4	11
M21	3	4	4	11
M22	3	4	4	11
M23	3	4	4	11
M24	3	4	4	11

Turbine code	December	January	February	Total
M25	3	4	4	11
M26	3	4	4	11
M27	3	4	4	11
M28	3	4	4	11
M29	3	4	4	11
M3	2	4	4	10
M30	3	4	4	11
M31	3	4	4	11
M32	3	4	4	11
M33	3	4	4	11
M34	3	4	4	11

Turbine code	December	January	February	Total
M35	3	4	4	11
M4	2	4	4	10
M5	2	4	4	10
M6	2	4	4	10
M7	2	4	4	10
M8	2	4	4	10
M9	2	4	4	10
VP1	2	4	4	10
VP2	2	4	4	10
ABZevs	2	4	4	10
Grand Total	253	480	456	1189

No body parts or intact remains of geese which could be considered as collision victims were detected after an accumulation of 1189 searches under 114 turbines in the period 01 December 2020 – 28 February 2021. Therefore, no evidence for collision of any goose species, including RBG, has been found in the winter 2020 – 2021 when geese were present, and turbines were operating.

There were circumstances in the 2020-2021 winter which required the Turbine Shutdown System (TSS). The number of TSS instances applied during this period are presented in Table 3.

Table 3. Number of Turbine Shutdown System applications in winter 2020-2021.

Date	Wind farm	Species	Number	Time stop	Time restart
31.01.2021	Wind Farm St. Nikola	<i>Pelicanus crispus</i>	3	12:04	12:28
08.02.2021	Wind Farm St. Nikola	<i>Haliaeetus albicilla</i>	1	11:48	12:15

Conclusions: wintering geese 2020-2021

The relatively mild 2020-2021 winter is probably the main reason for low number of observed geese of two species in ISPB territory.

Daily observations from December 2020 to February 2021 (inclusive) revealed that the recorded presence of geese in ISPB territory was compressed into a short time period within the winter, which was essentially the same as already established in 2018, 2019 and 2020 winter monitoring of the same territory as well as studies 2008 – 2018 in a part of the ISPB territory (SNWF).

The number of wintering geese observed in ISPB during winter broadly corresponds to the total number of wintering geese in the larger region of coastal Dobroudzha region; but is lower, because of relatively distant roosting sites to ISPB territory of wintering geese at the two fresh water lakes used for roosting – Durankulak and Shabla.

114 wind turbines covered by ISPB were not a source of collision mortality for wintering geese: an unusually small number flew through or fed within ISPB territory in 2020-21 winter. However, this same result has been found when considerably more geese utilised the ISPB territory (or a component: SNWF) in previous winters. The evidence for this is that no remains of geese that

could be attributed to collision with turbines were found during systematic searches under operational turbines: not only in the 2020-2021 winter but also in any of the 13 winters when all 114 turbines or 52 turbines at SNWF (part of ISPB) has been operational and searched systematically every winter season.

No displacement (disturbance) reaction from geese has been observed for the period 2008-2021 as a result of construction and operation of wind turbines in the ISPB territory. Observed numbers of geese of all species as well as observed spatial distribution of flying and feeding geese does not indicate gross displacement from the operational turbines or its immediate environs.

From research associated directly with ISPB described in the present and previous reports (and see previous SNWF winter reports on the AES Geo Energy website, and earlier surveys from this part of the same territory) the study area continues to be a feeding ground for RBG as well as GWFG, but it also remains an unimportant area for both species, as indicated in pre-construction studies. Consequently, and based on other studies, the investigated 114 wind turbines present no material threat through preventing use of food supplies: especially in light of other agricultural practices such as crop type and field size of the preferred crop of feeding geese.

Spring migration

During the spring monitoring, observations were made during all 61 days of the season (15 March -15 May), with registered migratory, soaring birds being detected over 70 % of the days in spring 2021. For the survey period, a total of 1012 migratory and resident birds were registered in spring 2021 (Table 4)

Table 4. Number of registered birds of all ecological groups by day during the spring migration in the territory covered by ISPB

Period	Number of birds in Spring 2021
15-31 March	590
1-30 April	354
1-15 May	68
Total for the period	1012




Figure 4. Dynamics of the spring migration of birds in the ISPB territory based on visual observations during the period 15 March - 15 May 2021

The most numerous birds in spring in the region for four migratory spring seasons were Common cormorants (*Phalacrocorax carbo*) and some birds of prey – Common buzzards (*Buteo buteo*), Red-footed falcon (*Falco vespertinus*), Common kestrels (*Falco tinnunculus*) and Marsh harriers (*Circus aeruginosus*). One of the most numerous species observed in spring seasons 2018 and 2019, the White pelican (*Pelecanus onocrotalus*), was not observed in the period of spring migration monitoring in 2020, but a flock of 120 White pelicans was observed 29 May 2020 in the ISPB territory. Only one White pelican was observed in 2021 during the period of spring migration monitoring in ISPB.

Table 5. Composition and number of registered bird species during the period 15 March - 15 May 2021 in the ISPB territory.

<i>Species name</i>	<i>Number of birds</i>	<i>Species name</i>	<i>Number of birds</i>
<i>A. cinerea</i>	58	<i>E. alba</i>	12
<i>A. heliaca</i>	1	<i>F. subbuteo</i>	5
<i>A. nisus</i>	13	<i>F. tinnunculus</i>	32
<i>A. pennata</i>	1	<i>F. vespertinus</i>	17
<i>A. pomarina</i>	3	<i>H. himantopus</i>	1
<i>A. purpurea</i>	22	<i>H. pennatus</i>	1
<i>B. buteo</i>	56	<i>L. ridibundus</i>	21
<i>B. rufinus</i>	30	<i>M. migrans</i>	2
<i>C. aeruginosus</i>	92	<i>N. nicticorax</i>	6
<i>C. ciconia</i>	24	<i>P. apricaria</i>	41
<i>C. corax</i>	16	<i>P. carbo</i>	469
<i>C. cyaneus</i>	4	<i>P. haliaetus</i>	1
<i>C. gallicus</i>	10	<i>P. onocrotalus</i>	1
<i>C. macrourus</i>	3	<i>T. tadorna</i>	63
<i>C. pygargus</i>	7	<i>Number of species</i>	29

Between 24 and 205 White storks (*Ciconia ciconia*) passed over the surveyed territory in the four spring seasons under the ISPB monitoring scheme. The European nesting population of the White stork is estimated to be between 180,000 and 220,000 pairs, with about 80 % of the species migrating along the wider western Black Sea region, which also covers a part of north-eastern Bulgaria. According to these values, White storks flying over the Kaliakra area, substantially east of the main migratory path of White storks along the western Black Sea migration corridor, were an insignificant proportion (between 0.02% and 0,09%) of the Via Pontica population. According to Shurulinkov et al. (2011), an estimate of the total population of White stork in SE Bulgaria flying along Via Pontica in spring was 23,358 individuals in their study period. In this respect our observations continue to confirm the low significance of the territory of Kaliakra as part of the migratory corridor for spring migrating White storks along the Via Pontica component of the larger flyway.

No stops of turbines were ordered under the Turbine Shutdown System (TSS) during the spring migration period of 2021. This was primarily because all the observed birds passing through the ISPB territory were outside the zone of the risk of collision with turbines.

In order to check the effectiveness of the ISPB to prevent collisions of spring migrating birds, each of the 114 turbines covered by the ISPB programme was checked at least once a week for collision victims. According to previously performed carcass removal and searcher efficiency tests during autumn migration and in winter at SNWF (and repeated in autumn 2018 for ISPB territory), this search regime of weekly searches provides for a cost-effective method, which can also be calibrated, to discover any bird strike fatalities which may be of concern. For details, see previous

studies of: <http://www.aesgeoenergy.com/site/Studies.html> and results of previous ISPB reports at: https://kaliakrabirdmonitoring.eu/Report_Autumn 2018.

Table 6. Number of turbines searched for collision victims in the territory of ISPB during the period 15 March-15 May 2021. The name of the wind farm operators and the number of the turbines used in the table: AE8/60 - AES Geo Energy Ltd., M1/35 - Kaliakra Wind Power, E1/8 - EVN Kavarna, DC1/2 - Degrets OOD, DBΓ1/5 - Disib OOD, DBΓ2MN600/DBΓ1HSW250 - Windex OOD, ABΓ4 - Long Man Invest OOD, ABBalgarevo - Long Man Energy OOD, AB3eac - Zevs Bonus OOD, VP1/2 - Vertikal-Petkov&Sie SD, ABΓ3 - Wind Park Kavarna East EOOD, ABΓ1/2 - Wind Park Kavarna West EOOD, AB Millennium Group Micon/ AB Millennium Group-Millennium Group OOD

Turbine	March	April	May	Total
ABBalgarevo	2	5	2	9
ABΓ1	9	11	2	22
ABΓ2	1	3	2	6
ABΓ3	1	3	2	6
ABΓ4	1	3	2	6
AB Millennium Group	2	5	2	9
AB Millennium Group Micon	2	5	2	9
AE10	2	5	2	9
AE11	2	5	2	9
AE12	3	4	2	9
AE13	3	4	2	9
AE14	2	4	2	8
AE15	2	4	2	8
AE16	2	5	2	9
AE17	2	5	2	9
AE18	3	4	2	9
AE19	3	4	2	9
AE20	2	4	2	8
AE21	2	5	2	9
AE22	2	5	2	9
AE23	2	5	2	9
AE24	2	4	3	9
AE25	2	4	3	9
AE26	2	5	2	9
AE27	2	5	2	9
AE28	2	5	2	9
AE29	2	4	3	9
AE31	3	4	2	9
AE32	3	4	2	9
AE33	3	4	2	9
AE34	3	4	2	9
AE35	3	4	2	9
AE36	2	4	2	8
AE37	3	4	2	9
AE38	2	4	2	8

Turbine	March	April	May	Total
AE39	2	4	2	8
AE40	2	4	3	9
AE41	2	4	3	9
AE42	2	4	3	9
AE43	2	4	3	9
AE44	2	4	3	9
AE45	2	5	2	9
AE46	3	4	2	9
AE47	3	4	2	9
AE48	3	4	2	9
AE49	3	4	2	9
AE50	3	4	2	9
AE51	3	4	2	9
AE52	3	4	2	9
AE53	3	4	2	9
AE54	3	4	2	9
AE55	3	4	2	9
AE56	3	4	2	9
AE57	3	4	2	9
AE58	3	4	2	9
AE59	3	4	2	9
AE60	3	4	2	9
AE8	2	4	2	8
AE9	2	4	2	8
DBΓ1	1	3	2	6
DBΓ1HSW250	2	4	3	9
DBΓ2	1	3	2	6
DBΓ2MN600	2	4	3	9
DBΓ3	1	3	2	6
DBΓ4	2	5	2	9
DBΓ5	2	5	2	9
DC1	2	5	2	9
DC2	2	5	2	9
E00	2	5	2	9
E01	2	4	3	9
E02	2	4	3	9

Turbine	March	April	May	Total
E04	2	4	3	9
E05	2	4	3	9
E07	2	4	3	9
E08	2	4	3	9
E09	2	5	2	9
M1	2	5	2	9
M10	2	5	2	9
M11	2	5	2	9
M12	3	4	2	9
M13	3	4	2	9
M14	3	4	2	9
M15	3	4	2	9
M16	3	4	2	9
M17	3	4	2	9
M18	3	4	2	9
M19	3	4	2	9
M2	2	5	2	9
M20	3	4	2	9
M21	3	4	2	9
M22	3	4	2	9
M23	3	4	2	9
M24	3	4	2	9

Turbine	March	April	May	Total
M25	3	4	2	9
M26	3	4	2	9
M27	3	4	2	9
M28	3	4	2	9
M29	3	4	2	9
M3	2	5	2	9
M30	3	4	2	9
M31	3	4	2	9
M32	3	4	2	9
M33	3	4	2	9
M34	3	4	2	9
M35	3	4	2	9
M4	2	5	2	9
M5	2	5	2	9
M6	2	5	2	9
M7	2	5	2	9
M8	2	5	2	9
M9	2	5	2	9
VP1	2	5	2	9
VP2	2	5	2	9
AB3eBc	1	3	2	6
Grand Total	277	489	244	1010

13 records of dead birds after collision with wind turbines were documented during the 2021 spring migration of birds in ISPB territory (Table 7). Most numerous among the confirmed collision victims during the study period were Skylarks (*Alauda arvensis*) and Calandra lark (*Melanocorypha calandra*). No case of collision with the turbines of a target bird species for the period of TSS application in ISPB was registered during the monitoring in spring 2021 (the target species are listed at <https://kaliakrabirdmonitoring.eu/>).

Table 7. Confirmed collision victims and species' conservation status as recorded during the 2021 spring migration period.

English name	Species name	Number of birds	Red Data Book	IUCN
Calandra lark	<i>Melanocorypha calandra</i>	4	Endangered	LC
Common starling	<i>Sturnus vulgaris</i>	1	Not listed	LC
Common pheasant	<i>Phasianus colchicus</i>	1	Not listed	LC
Corn bunting	<i>Emberiza calandra</i>	1	Not listed	LC
Grey partridge	<i>Perdix perdix</i>	1	Not listed	LC
Skylark	<i>Alauda arvensis</i>	3	Not listed	LC
Willow warbler	<i>Phylloscopus trochilus</i>	1	Not listed	LC
Yellow legged gull	<i>Larus michahellis</i>	1	Not listed	LC

Conclusions: spring migration

During the monitoring, there were no apparent changes in the main characteristics of the ornithofauna typical for the spring migration in the whole country and the specific characteristics of the species composition and phenology of spring bird migration in NE Bulgaria.

The results of the monitoring confirmed the relatively low importance of the ISPB territory for migratory birds in spring and the absence of negative influence of the operating wind farms on bird populations during their spring migration.

During the migration periods, the species composition, the dynamics in number of birds, the daily activity, the height of the flights, as well as the feeding, resting and roost sites of the flying birds passing through the area indicated the absence of a barrier effect of the 114 wind turbines.

The data presented in this report confirmed the absence of any adverse impact on sensitive bird species of the orders Ciconiiformes, Pelecaniformes, Falconiformes, Gruiformes using migratory ascending air flows (thermals) for movement over long distances.

All these species were found to occasionally cross the study site, and their observed behaviour in respect to wind turbines did not indicate major changes which would impact on the energetics of these species during daily movements.

The quantitative characteristics of bird migration in the ISPB area during spring 2021, and the absence of mortality among the target bird species allows a continued conclusion that the studied wind farms do not present a risk of adverse impact to migratory birds. The application of the ISPB’s safeguards potentially was and can be an ongoing contributory part of the minimal risk posed to birds from wind farms in the Kaliakra region.

Autumn migration

During the autumn monitoring, observations were made during all 92 days of the season 2021 (01.08-31.10.2021).




Figure 5. Number of registered birds by months during the autumn migration in the territory of ISPB.




Figure 6. Dynamics of the autumn migration of the flying bird species in the ISPB territory according to visual observations during the period 01 August - 31 October 2021. Letters indicate the direction of wind in days with increased number of migrating birds.

The number of birds in the ISPB study area apparently depended on the direction of the wind in autumn 2021 (as in previous autumns). Of the seven peak days with intense migratory flights of birds: in three, west winds prevailed; in two, north-west and in another two days, the wind direction was north.(Figure 6).

The monitoring from 1 August to 31 October 2021 recorded 22983 individual birds, assigned to 46 bird species. The numbers of individuals recorded by species during autumn migration in 2021 are shown in Table 8.

Table 8. Composition of species and number of registered birds over the period 01 August to 31 October 2021 in the ISPB territory.

Species name	Number
<i>A. alba</i>	5
<i>A. brevipes</i>	194
<i>A. gentilis</i>	1
<i>A. nisus</i>	150
<i>A. cinerea</i>	49
<i>A. purpurea</i>	1
<i>A. pennata</i>	17
<i>A. pomarina</i>	27
<i>A. heliaca</i>	1
<i>B. buteo</i>	615
<i>B. rufinus</i>	8
<i>C. aeruginosus</i>	202
<i>C. cyaneus</i>	18
<i>C. pygargus</i>	27
<i>C. macrourus</i>	6
<i>C. gallicus</i>	63
<i>C. ciconia</i>	12859
<i>C. nigra</i>	17
<i>C. garrulus</i>	14
<i>C. corax</i>	13
<i>C. coturnix</i>	1
<i>C. olor</i>	5
<i>C. palumbus</i>	500

Species name	Number
<i>F. vespertinus</i>	397
<i>F. subbuteo</i>	34
<i>F. peregrinus</i>	3
<i>F. tinnunculus</i>	94
<i>M. migrans</i>	28
<i>M. milvus</i>	1
<i>M. apiaster</i>	2374
<i>G. grus</i>	251
<i>G. virgo</i>	1
<i>G. fulvus</i>	1
<i>L. michahellis</i>	626
<i>L. melanocephalus</i>	450
<i>L. fuscus</i>	1
<i>L. ridibundus</i>	35
<i>H. minutus</i>	45
<i>H. rustica</i>	200
<i>P. carbo</i>	319
<i>P. onocrotalus</i>	1449
<i>P. crispus</i>	2
<i>P. apivorus</i>	1852
<i>P. haliaetus</i>	8
<i>R. riparia</i>	17
<i>U. epops</i>	2

The most numerous migrating birds recorded in autumn 2021 were white storks (*Ciconia ciconia*) with over 12,000 individuals registered. Within the other soaring species, the most numerous recorded birds involved honey buzzards (*P. apivorus*) and great white pelicans (*P. onocrotalus*) with over 1400 individuals of each species (Table 8). Seven new species were recorded in autumn 2021. The newly observed species were great egret (*Ardea alba*), common quail (*Coturnix coturnix*), mute swan (*Cygnus olor*), Mediterranean gull (*Larus melanocephalus*), black-headed gull (*Larus ridibundus*), little gull (*Hydrocoloeus minutus*), and Eurasian hoopoe (*Upupa epops*). The hoopoe is a common bird species and the fact it was not registered in previous autumn seasons is probably related to the habitats around the constant observation points which are located in agrarian fields.

As a result of the simultaneous observations at five constant observation points and three radar systems (Figure 1) during the whole period of the 2021 autumn migration, there were five stops of four groups of turbines and one complete wind farm (KWP) in the territory of the Kaliakra SPA and adjacent territories. The stop orders given to the engineers on duty were executed in a timely manner, thus avoiding any collision risk of bird passing through the territory. Detailed information on the duration of the ordered stops is given in Table 9.

Table 9. Data for stops of wind turbines ordered by field observers during the autumn migration of birds 2021.

Date	Wind Farm	Turbine code №/ Group	Species	Number of birds	Time stop	Time restart
02.10.2021	SNWF	A zone	<i>P. onocrotalus</i> <i>G. grus</i>	450 13	15:42:00	15:48:00
10.10.2021	KWP	-	<i>P. onocrotalus</i>	35	13:08:00	13:25:00
11.10.2021	SNWF	A zone	<i>G. fulvus</i>	1	10:28:00	10:33:00
11.10.2021	SNWF	F zone	<i>G. fulvus</i>	1	10:45:00	10:52:00

According to additional previously performed carcass removal and searcher efficiency tests during autumn migration and in winter at SNWF, a weekly search regime provides for a cost-effective method, which can also be calibrated, to discover any bird strike fatalities which may be of concern. Hence a frequency of four searches per month under every turbine allows estimation of the mortality of birds from collision with the turbines in the ISPB. This allows estimation of bird mortality from collision with the turbines in the Kaliakra SPA and others of the total 114 wind turbines included in the ISPB. For details of relevant previous studies at SNWF within the wider ISPB territory, see: <http://www.aesgeoenergy.com/site/Studies.html>

Table 10. Number of turbines searched for collision victims in the territory of ISPB during the period 01 August to 31 October 2021. The name of the wind farm operators and the number of the turbines used in the table: AE8/60 - AES Geo Energy Ltd., M1/35 - Kaliakra Wind Power, E1/8 - EVN Kavarna, DC1/2 - Degrets OOD, DBГ1/5 - Disib OOD, DBГ2MN600/DBГ1HSW250 - Windex OOD, ABГ4 - Long Man Invest OOD, ABBalgarevo - Long Man Energy OOD, AB3eac - Zevs Bonus OOD, VP1/2 - Vertikal-Petkov&Sie SD, ABГ3 - Wind Park Kavarna East EOOD, ABГ1/2 - Wind Park Kavarna West EOOD, AB Millennium group Micon/ AB Millennium group - Millennium Group OOD.

Turbine	Aug.	Sep.	Oct.	Total
ABBalgarevo	3	4	4	11
ABГ1	3	4	4	11
ABГ2	3	4	4	11
ABГ3	3	4	4	11
ABГ4	3	4	4	11
ABMillenium group	3	4	4	11
ABMillenium group Micon	3	4	2	9

Turbine	Aug.	Sep.	Oct.	Total
AE10	3	4	4	11
AE11	3	4	4	11
AE12	4	4	4	12
AE13	4	6	4	14
AE14	3	4	4	11
AE15	3	4	4	11
AE16	3	4	4	11

Turbine	Aug.	Sep.	Oct.	Total
AE17	3	4	4	11
AE18	4	4	4	12
AE19	4	4	4	12
AE20	3	4	4	11
AE21	3	4	4	11
AE22	3	4	4	11
AE23	3	4	4	11
AE24	3	4	4	11
AE25	3	4	4	11
AE26	3	4	4	11
AE27	3	4	3	10
AE28	3	4	3	10
AE29	3	4	4	11
AE31	4	6	4	14
AE32	4	6	4	14
AE33	4	6	4	14
AE34	4	6	4	14
AE35	4	6	4	14
AE36	3	4	4	11
AE37	4	4	4	12
AE38	3	4	4	11
AE39	3	4	4	11
AE40	3	4	4	11
AE41	3	4	4	11
AE42	3	4	4	11
AE43	3	4	4	11
AE44	3	4	4	11
AE45	3	4	3	10
AE46	4	4	4	12
AE47	4	4	4	12
AE48	4	4	4	12
AE49	4	4	4	12
AE50	4	6	4	14
AE51	4	4	4	12
AE52	4	4	4	12
AE53	4	4	4	12
AE54	4	4	4	12
AE55	4	4	4	12
AE56	4	4	4	12
AE57	4	4	4	12
AE58	4	4	4	12
AE59	4	4	4	12
AE60	4	6	4	14
AE8	3	4	4	11
AE9	3	4	4	11
DBG1	3	4	4	11

Turbine	Aug.	Sep.	Oct.	Total
DBG1HSW250	2	4	4	10
DBG2	3	4	4	11
DBG2MN600	2	4	4	10
DBG3	3	4	4	11
DBG4	3	4	3	10
DBG5	3	4	3	10
DC1	3	4	3	10
DC2	3	4	3	10
E00	3	4	4	11
E01	2	4	4	10
E02	2	4	4	10
E04	2	4	4	10
E05	2	4	4	10
E07	2	4	4	10
E08	2	4	4	10
E09	3	4	4	11
M1	3	4	4	11
M10	3	5	3	11
M11	3	5	3	11
M12	2	4	3	9
M13	2	4	3	9
M14	2	4	3	9
M15	2	4	3	9
M16	2	4	3	9
M17	2	4	3	9
M18	2	4	3	9
M19	2	4	3	9
M2	3	4	4	11
M20	2	5	4	11
M21	2	5	4	11
M22	2	5	4	11
M23	2	5	4	11
M24	2	5	4	11
M25	2	5	4	11
M26	2	5	4	11
M27	2	5	4	11
M28	4	4	4	12
M29	4	4	4	12
M3	3	4	4	11
M30	4	4	4	12
M31	4	4	4	12
M32	4	4	4	12
M33	4	4	4	12
M34	4	4	4	12
M35	4	4	4	12
M4	3	5	3	11

Turbine	Aug.	Sep.	Oct.	Total
M5	3	5	3	11
M6	3	5	3	11
M7	3	5	3	11
M8	3	5	3	11
M9	3	5	3	11

Turbine	Aug.	Sep.	Oct.	Total
VP1	3	4	4	11
VP2	3	4	4	11
ABZevs	3	4	4	11
Grand Total	351	488	431	1270

As a result of 1270 single inspections of 114 individual turbines between 1 August and 31 October 2021, a total of 11 dead birds of nine species were identified. The number of identified collision victims by species are given in Table 11.

Table 11. Victims of collision with turbines during the autumn migration period in 2021 according to the Red Data Book for Bulgaria and IUCN conservation status classifications (LC = Least Concern)

Species name	Scientific name	Number	Red Data Book	IUCN
<i>Calandra lark</i>	<i>Melanocorypha calandra</i>	1	endangered	LC
<i>Corn bunting</i>	<i>Emberiza calandra</i>	2	not listed	LC
<i>Domestic pigeon</i>	<i>Columba livia domestica</i>	1	not listed	LC
<i>Eurasian blackcap</i>	<i>Sylvia atricapilla</i>	1	not listed	LC
<i>European Bee-eater</i>	<i>Merops apiaster</i>	1	not listed	LC
<i>Lesser grey shrike</i>	<i>Lanius minor</i>	1	not listed	LC
<i>Red-backed shrike</i>	<i>Lanius collurio</i>	2	not listed	LC
<i>Thrush nightingale</i>	<i>Luscinia luscinia</i>	1	not listed	LC
<i>Yellow-legged gull</i>	<i>Larus michahellis</i>	1	not listed	LC

Eight of the bird species identified as victims are not listed in the Red Data Book of Bulgaria. Calandra lark is listed in Bulgarian Red Data Book but in the period of autumn bird migration all birds found during carcass searches are migrants and have to be considered as immigrants into Bulgaria. Therefore, for the evaluation of the population level impact of the additive mortality of wind turbines included in the monitoring, the international bird species status must be applied. IUCN classifications as Least Concern (LC) were appropriate to all species identified as collision victims. The category Least Concern indicates that the species has been evaluated against the Red List criteria and does not qualify for Critically Endangered, Endangered, Vulnerable or Near Threatened. Widespread and abundant taxa are included in this category. All recorded victims were not among the target ISPB species.

Conclusions: autumn migration

During the monitoring of ISPB territory, there were no substantive differences in the main characteristics of the ornithofauna typical for the autumn migration in the whole country and the specific characteristics of species' composition and phenology of bird migration in NE Bulgaria.

The results of the monitoring confirmed the relatively low importance of the ISPB territory for the birds flying through or over it and no apparent negative influence of the operating wind farms on bird populations during their autumn migration.

The migration periods, the species composition, the dynamics in number of birds, the daily activity, the height of the flights, as well as the feeding, resting and roost sites of the flying birds passing through the area and the observation points indicated the absence of a barrier effect of the 114 wind turbines covered by ISPB in autumn migration period.

The data presented in this report confirmed the absence of impact on sensitive bird species using migratory upward airflows (thermals) to move (soaring) over long distances in autumn migration period.

All these species were found during the study to cross the site using suitable habitats without the need to increase their energy losses in their daily movements and to change their migratory strategy in the autumn period.

The quantitative characteristics of bird migration in the ISPB area during autumns of 2018, 2019, 2020 and 2021 and the absence of mortality among the target bird species allows a continued conclusion that the studied wind farms do not present a risk of adverse impact to migratory birds. The application of the ISPB's safeguards potentially was and can be an ongoing contributory part of the minimal risk posed to birds from wind farms in the Kaliakra region.

List of participants in the observations

Ø Prof. Dr Pavel Zehtindjiev – Senior field ornithologist

More than 25 years of research experience in ornithology. Author of more than 85 scientific publications in international journals with an impact on the scientific field of bird biology, ecology and ecosystem conservation. Member of the European Ornithological Union and many nature conservation organizations. Winner of the Revolutionary Discovery Award for the Ornithology of the American Ornithological Society for 2016 - The Cooper Ornithological Society.

Over 10 years of experience in impact monitoring study of wind turbines in the study area.

Ø Dr Viktor Vasilev – Field ornithologist

Senior researcher in the Faculty of Biology, University of Shumen.

Member of BSPB and participant in number of conservation projects in Bulgaria.

Author of over 20 scientific publications in international journals. Member of BSPB.

Ø Veselina Raikova - Field ornithologist

Natural History Museum of Varna. Member of BSPB. Author of more than 10 publications in international scientific journals. Over 10 years of experience in impact monitoring study of wind turbines in the study area.

Ø Ivaylo Raykov - Field ornithologist

Museum of Natural History, Varna. Member of BSPB. Author of over 20 scientific publications in international journals.

Five years of experience in impact monitoring in the region of Kaliakra.

Ø Kiril Bedev - Field ornithologist

Researcher in Institute of Biodiversity and Ecosystem Research at the Bulgarian Academy of Sciences.

Active member of conservation organization Green Balkans. Long term study on migrating birds and biodiversity of Burgas lakes. Author of three articles in Bulgarian Red Data Book. Expertise in biotechnology, conservation biology and environmental monitoring. Over seven years of experience in impact monitoring of wind parks in Bulgaria. Member of Balkani NGO for conservation of birds and nature.

Ø Janko Jankov - Field ornithologist

Student in Biology, University of Shumen. Over seven years of experience in impact monitoring of birds in Wind Park projects in NE Bulgaria. Member of BSPB.

Ø Nikolay Velichkov - Field ornithologist

Field studies of the distribution and number of breeding bird species ENVEKO, Inspection of use of pesticides and pedigrees in the framework of the project "Urgent measures for the protection of the Egyptian Vulture (*Neophron percnopterus*) BSPB".

Monitoring the migration of birds species composition and the number of nesting fauna 2007-2012 "Ecotan" EOOD. 10 years of experience in impact monitoring study of wind turbines in the study area

Ø Svetoslav Stoyanov - Field ornithologist

Bachelor in Biology diploma from Shumen University. Participant in numerous conservation projects of BSPB – BirdLife Bulgaria. Midwinter counts of waterfowl birds in Bulgaria and white stork census expert. Monitoring the migration of birds species composition and the number of nesting fauna 2007-2012 "Ecotan" EOOD. Over 10 years of experience in impact monitoring study of wind turbines in the study area

Ø Minko Madjarov - Field ornithologist

Experienced ornithologist with over 10 years of field work in conservation projects of BSPB (BirdLife Bulgaria). Participant in the project - Mapping and Determination of the Conservation Status of Natural Habitats and Species - Phase 1, Lot 7 - Determination and Minimization of Risks for Wild Birds. Union Econet – MOEW. Birdwatching guide for over 10 years.

Ø Jelyazko Dimitrov - Field ornithologist

Member of BSPB from 31.12.2006 to 31.12.2010. Trained to monitor the severity of collisions of birds with wind turbines.

Ø Dimitar Dimitrov - Field ornithologist

Student in Biology at Sofia University Kliment Ohridski. Field activities - participation in a number of field studies - monitoring of some important zones on the territory of Bulgaria. (Durankulak lake and the Shabla lake complex (2010 - 2013) and the Soil Field (2014-2017), regular winter monitoring of waterfowl in Shabla and Durankulak Lake in connection with the Life + project (2011 - 2017), monitoring of *Spermophilus cittelus* in the reintroduced colony near Kotel (2017), census of cetacean mammals on the northern Black Sea coast with ECO-Nord association, voluntary eye initiatives on reintroduction of the griffon vulture in the Kresna Gorge.

Ø Boyan Michev - Field ornithologist

PhD student at the Institute of Biodiversity and Ecosystem Research - BAS. He works in Risk Assessment and Conservation Biology department. Expert in the use of radars to study bird migration. Member of the European Migration Tracking Network through meteorological radars.