

Appendix D

Tables of predicted relative abundance as a proportion of total relative abundance in the study area by BOEM wind energy planning/lease area and season for each species modeled

List of Tables

Arctic Tern (ARTE) proportional relative abundance by BOEM area and season.....	2
Atlantic Puffin (ATPU) proportional relative abundance by BOEM area and season.....	3
Audubon’s Shearwater (AUSH) proportional relative abundance by BOEM area and season.....	4
Black-capped Petrel (BCPE) proportional relative abundance by BOEM area and season.....	5
Black Guillemot (BLGU) proportional relative abundance by BOEM area and season.....	6
Black-legged Kittiwake (BLKI) proportional relative abundance by BOEM area and season.....	7
Black Scoter (BLSC) proportional relative abundance by BOEM area and season.....	8
Bonaparte’s Gull (BOGU) proportional relative abundance by BOEM area and season.....	9
Brown Pelican (BRPE) proportional relative abundance by BOEM area and season.....	10
Band-rumped Storm-Petrel (BRSP) proportional relative abundance by BOEM area and season.....	11
Bridled Tern (BRTE) proportional relative abundance by BOEM area and season.....	12
Common Eider (COEI) proportional relative abundance by BOEM area and season.....	13
Common Loon (COLO) proportional relative abundance by BOEM area and season.....	14
Common Murre (COMU) proportional relative abundance by BOEM area and season.....	15
Cory’s Shearwater (COSH) proportional relative abundance by BOEM area and season.....	16
Common Tern (COTE) proportional relative abundance by BOEM area and season.....	17
Double-crested Cormorant (DCCO) proportional relative abundance by BOEM area and season.....	18
Dovekie (DOVE) proportional relative abundance by BOEM area and season.....	19
Great Black-backed Gull (GBBG) proportional relative abundance by BOEM area and season.....	20
Great Shearwater (GRSH) proportional relative abundance by BOEM area and season.....	21
Great Skua (GRSK) proportional relative abundance by BOEM area and season.....	22
Herring Gull (HERG) proportional relative abundance by BOEM area and season.....	23
Horned Grebe (HOGR) proportional relative abundance by BOEM area and season.....	24
Laughing Gull (LAGU) proportional relative abundance by BOEM area and season.....	25
Leach’s Storm-Petrel (LESP) proportional relative abundance by BOEM area and season.....	26
Least Tern (LETE) proportional relative abundance by BOEM area and season.....	27
Long-tailed Duck (LTDU) proportional relative abundance by BOEM area and season.....	28
Manx Shearwater (MASH) proportional relative abundance by BOEM area and season.....	29
Northern Fulmar (NOFU) proportional relative abundance by BOEM area and season.....	30
Northern Gannet (NOGA) proportional relative abundance by BOEM area and season.....	31
Parasitic Jaeger (PAJA) proportional relative abundance by BOEM area and season.....	32
Pomarine Jaeger (POJA) proportional relative abundance by BOEM area and season.....	33
Razorbill (RAZO) proportional relative abundance by BOEM area and season.....	34
Ring-billed Gull (RBGU) proportional relative abundance by BOEM area and season.....	35
Red-breasted Merganser (RBME) proportional relative abundance by BOEM area and season.....	36
Red Phalarope (REPH) proportional relative abundance by BOEM area and season.....	37
Red-necked Phalarope (RNPH) proportional relative abundance by BOEM area and season.....	38
Roseate Tern (ROST) proportional relative abundance by BOEM area and season.....	39
Royal Tern (ROYT) proportional relative abundance by BOEM area and season.....	40
Red-throated Loon (RTLO) proportional relative abundance by BOEM area and season.....	41
Sooty Shearwater (SOSH) proportional relative abundance by BOEM area and season.....	42
Sooty Tern (SOTE) proportional relative abundance by BOEM area and season.....	43
South Polar Skua (SPSK) proportional relative abundance by BOEM area and season.....	44
Surf Scoter (SUSC) proportional relative abundance by BOEM area and season.....	45
Thick-billed Murre (TBMU) proportional relative abundance by BOEM area and season.....	46
Wilson’s Storm-Petrel (WISP) proportional relative abundance by BOEM area and season.....	47
White-winged Scoter (WWSC) proportional relative abundance by BOEM area and season.....	48




Figure D1. BOEM Wind Energy Lease/Planning Areas.

Predicted relative abundance of ARTE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478		3.0e-05		
Rhode Island / Massachusetts OCS-A 0486		9.9e-05		
Rhode Island / Massachusetts OCS-A 0487		6.8e-05		
Massachusetts OCS-A 0500		1.9e-04		
Massachusetts OCS-A 0501		1.8e-04		
Massachusetts OCS-A 0502		2.7e-04		
Massachusetts OCS-A 0503		1.8e-04		
New York Proposed Commercial Lease - Unsolicited		3.9e-05		
New York OCS-A 0512		7.4e-05		
New Jersey OCS-A 0499		1.6e-04		
New Jersey OCS-A 0498		1.4e-04		
Delaware OCS-A 0482		8.4e-05		
Maryland OCS-A 0489		2.8e-05		
Maryland OCS-A 0490		4.1e-05		
Virginia OCS-A 0483		9.7e-05		
Virginia OCS-A 0497		2.4e-06		
North Carolina OCS-A 0508		1.1e-04		
North Carolina WEA - Wilmington West		4.3e-05		
North Carolina WEA - Wilmington East		1.1e-04		
South Carolina Call Area - Grand Strand		5.2e-04		
South Carolina Call Area - Cape Romain		1.3e-04		
South Carolina Call Area - Winyah		2.9e-05		
South Carolina Call Area - Charleston		2.9e-05		
All		2.7e-03		

Predicted relative abundance of ATPU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.6e-05	1.9e-05	2.5e-05	1.1e-05
Rhode Island / Massachusetts OCS-A 0486	1.1e-04	5.4e-05	9.4e-05	5.8e-05
Rhode Island / Massachusetts OCS-A 0487	9.7e-05	3.6e-05	6.2e-05	6.4e-05
Massachusetts OCS-A 0500	3.2e-04	1.0e-04	1.8e-04	2.3e-04
Massachusetts OCS-A 0501	2.2e-04	9.3e-05	1.5e-04	2.2e-04
Massachusetts OCS-A 0502	3.6e-04	1.3e-04	2.1e-04	4.0e-04
Massachusetts OCS-A 0503	2.3e-04	7.5e-05	1.2e-04	2.5e-04
New York Proposed Commercial Lease - Unsolicited	3.0e-05	2.0e-05	3.3e-05	3.6e-05
New York OCS-A 0512	4.2e-05	3.9e-05	7.2e-05	5.2e-05
New Jersey OCS-A 0499	8.5e-05	7.9e-05	1.4e-04	7.4e-05
New Jersey OCS-A 0498	7.1e-05	6.8e-05	1.2e-04	6.6e-05
Delaware OCS-A 0482	4.1e-05	4.0e-05	7.3e-05	3.9e-05
Maryland OCS-A 0489	1.4e-05	1.3e-05	2.5e-05	1.1e-05
Maryland OCS-A 0490	2.0e-05	1.9e-05	3.5e-05	2.5e-05
Virginia OCS-A 0483	4.6e-05	4.0e-05	8.0e-05	6.7e-05
Virginia OCS-A 0497	1.1e-06	1.0e-06	2.0e-06	1.2e-06
North Carolina OCS-A 0508	6.5e-05	3.9e-05	8.4e-05	1.3e-04
North Carolina WEA - Wilmington West	1.2e-05	2.0e-05	4.1e-05	1.2e-05
North Carolina WEA - Wilmington East	4.1e-05	4.7e-05	1.1e-04	4.7e-05
South Carolina Call Area - Grand Strand	1.5e-04	2.5e-04	5.1e-04	1.5e-04
South Carolina Call Area - Cape Romain	3.5e-05	6.1e-05	1.3e-04	2.9e-05
South Carolina Call Area - Winyah	2.2e-05	1.3e-05	2.9e-05	1.4e-05
South Carolina Call Area - Charleston	9.9e-06	1.5e-05	3.2e-05	1.1e-05
All	2.0e-03	1.3e-03	2.4e-03	2.0e-03

Predicted relative abundance of AUSH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.6e-06	2.6e-06	2.5e-06	1.2e-09
Rhode Island / Massachusetts OCS-A 0486	5.2e-06	9.5e-06	8.9e-06	3.8e-09
Rhode Island / Massachusetts OCS-A 0487	3.6e-06	7.1e-06	6.1e-06	2.6e-09
Massachusetts OCS-A 0500	1.0e-05	1.9e-05	1.7e-05	7.3e-09
Massachusetts OCS-A 0501	8.9e-06	1.7e-05	1.5e-05	6.5e-09
Massachusetts OCS-A 0502	1.3e-05	3.3e-05	2.2e-05	9.6e-09
Massachusetts OCS-A 0503	7.5e-06	2.2e-05	1.2e-05	5.4e-09
New York Proposed Commercial Lease - Unsolicited	2.2e-06	4.0e-06	4.0e-06	1.6e-09
New York OCS-A 0512	4.2e-06	7.2e-06	8.2e-06	3.2e-09
New Jersey OCS-A 0499	9.9e-06	1.7e-05	2.0e-05	7.2e-09
New Jersey OCS-A 0498	8.7e-06	1.5e-05	1.8e-05	6.3e-09
Delaware OCS-A 0482	5.2e-06	9.2e-06	1.1e-05	3.8e-09
Maryland OCS-A 0489	1.8e-06	3.2e-06	3.6e-06	1.3e-09
Maryland OCS-A 0490	2.6e-06	4.6e-06	6.2e-06	1.9e-09
Virginia OCS-A 0483	6.4e-06	1.5e-05	2.2e-05	4.5e-09
Virginia OCS-A 0497	1.6e-07	3.3e-07	4.0e-07	1.1e-10
North Carolina OCS-A 0508	6.9e-06	2.9e-05	4.3e-05	5.1e-09
North Carolina WEA - Wilmington West	2.9e-06	8.6e-06	6.6e-06	1.2e-06
North Carolina WEA - Wilmington East	1.2e-05	1.0e-04	5.5e-05	7.6e-07
South Carolina Call Area - Grand Strand	3.6e-05	1.1e-04	9.6e-05	6.9e-07
South Carolina Call Area - Cape Romain	9.0e-06	3.5e-05	2.1e-05	8.1e-09
South Carolina Call Area - Winyah	8.8e-06	7.9e-05	4.8e-04	2.7e-07
South Carolina Call Area - Charleston	5.0e-06	2.1e-05	2.6e-05	2.1e-08
All	1.7e-04	5.7e-04	9.1e-04	3.0e-06

Predicted relative abundance of BCPE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	6.1e-08	3.8e-10	8.8e-09	5.1e-11
Rhode Island / Massachusetts OCS-A 0486	2.0e-07	1.3e-09	2.9e-08	1.7e-10
Rhode Island / Massachusetts OCS-A 0487	1.4e-07	8.9e-10	2.0e-08	1.2e-10
Massachusetts OCS-A 0500	3.9e-07	2.4e-09	5.6e-08	3.2e-10
Massachusetts OCS-A 0501	3.5e-07	2.2e-09	5.0e-08	2.9e-10
Massachusetts OCS-A 0502	5.3e-07	3.2e-09	7.4e-08	4.3e-10
Massachusetts OCS-A 0503	3.0e-07	1.8e-09	4.2e-08	2.4e-10
New York Proposed Commercial Lease - Unsolicited	8.4e-08	5.3e-10	1.2e-08	7.1e-11
New York OCS-A 0512	1.6e-07	1.0e-09	2.4e-08	1.4e-10
New Jersey OCS-A 0499	3.8e-07	2.4e-09	5.5e-08	3.2e-10
New Jersey OCS-A 0498	3.4e-07	2.1e-09	4.8e-08	2.8e-10
Delaware OCS-A 0482	2.0e-07	1.3e-09	2.9e-08	1.7e-10
Maryland OCS-A 0489	6.9e-08	4.3e-10	9.8e-09	5.7e-11
Maryland OCS-A 0490	1.0e-07	6.2e-10	1.4e-08	8.1e-11
Virginia OCS-A 0483	2.5e-07	1.8e-09	3.4e-08	2.0e-10
Virginia OCS-A 0497	6.1e-09	4.3e-11	8.4e-10	4.8e-12
North Carolina OCS-A 0508	2.8e-07	2.0e-09	3.7e-08	2.1e-10
North Carolina WEA - Wilmington West	1.2e-07	7.3e-10	1.5e-08	9.0e-11
North Carolina WEA - Wilmington East	3.4e-07	5.5e-08	4.0e-08	2.6e-10
South Carolina Call Area - Grand Strand	1.4e-06	9.5e-09	1.9e-07	1.1e-09
South Carolina Call Area - Cape Romain	3.6e-07	7.2e-09	4.7e-08	2.7e-10
South Carolina Call Area - Winyah	2.7e-07	1.1e-08	1.2e-08	6.7e-11
South Carolina Call Area - Charleston	1.1e-07	4.6e-09	1.1e-08	6.2e-11
All	6.5e-06	1.1e-07	8.6e-07	5.0e-09

Predicted relative abundance of BLGU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478		8.0e-05		
Rhode Island / Massachusetts OCS-A 0486		2.2e-04		
Rhode Island / Massachusetts OCS-A 0487		1.5e-04		
Massachusetts OCS-A 0500		4.0e-04		
Massachusetts OCS-A 0501		3.6e-04		
Massachusetts OCS-A 0502		5.0e-04		
Massachusetts OCS-A 0503		2.7e-04		
New York Proposed Commercial Lease - Unsolicited		8.4e-05		
New York OCS-A 0512		1.7e-04		
New Jersey OCS-A 0499		3.5e-04		
New Jersey OCS-A 0498		3.0e-04		
Delaware OCS-A 0482		1.8e-04		
Maryland OCS-A 0489		5.9e-05		
Maryland OCS-A 0490		8.3e-05		
Virginia OCS-A 0483		1.8e-04		
Virginia OCS-A 0497		4.6e-06		
North Carolina OCS-A 0508		1.8e-04		
North Carolina WEA - Wilmington West		9.2e-05		
North Carolina WEA - Wilmington East		2.2e-04		
South Carolina Call Area - Grand Strand		1.2e-03		
South Carolina Call Area - Cape Romain		3.2e-04		
South Carolina Call Area - Winyah		6.1e-05		
South Carolina Call Area - Charleston		7.0e-05		
All		5.5e-03		

Predicted relative abundance of BLKI as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	6.4e-05		2.0e-04	1.3e-04
Rhode Island / Massachusetts OCS-A 0486	3.6e-04		9.0e-04	3.9e-04
Rhode Island / Massachusetts OCS-A 0487	3.7e-04		4.4e-04	2.9e-04
Massachusetts OCS-A 0500	1.1e-03		1.0e-03	8.1e-04
Massachusetts OCS-A 0501	1.1e-03		8.9e-04	7.3e-04
Massachusetts OCS-A 0502	1.9e-03		1.1e-03	1.0e-03
Massachusetts OCS-A 0503	1.3e-03		5.8e-04	5.8e-04
New York Proposed Commercial Lease - Unsolicited	2.4e-04		1.8e-04	1.5e-04
New York OCS-A 0512	2.4e-04		3.7e-04	2.7e-04
New Jersey OCS-A 0499	2.0e-04		4.5e-04	3.9e-04
New Jersey OCS-A 0498	1.6e-04		3.5e-04	2.7e-04
Delaware OCS-A 0482	8.5e-05		2.0e-04	8.9e-05
Maryland OCS-A 0489	2.8e-05		6.2e-05	2.9e-05
Maryland OCS-A 0490	4.0e-05		9.0e-05	4.4e-05
Virginia OCS-A 0483	8.4e-05		1.6e-04	9.1e-05
Virginia OCS-A 0497	1.9e-06		4.0e-06	2.1e-06
North Carolina OCS-A 0508	9.8e-05		1.3e-04	1.6e-04
North Carolina WEA - Wilmington West	2.2e-05		6.4e-05	4.5e-05
North Carolina WEA - Wilmington East	6.7e-05		1.6e-04	1.7e-04
South Carolina Call Area - Grand Strand	2.8e-04		7.7e-04	4.6e-04
South Carolina Call Area - Cape Romain	6.9e-05		2.0e-04	1.1e-04
South Carolina Call Area - Winyah	1.7e-05		2.9e-05	2.0e-05
South Carolina Call Area - Charleston	2.0e-05		4.2e-05	2.6e-05
All	7.8e-03		8.4e-03	6.3e-03

Predicted relative abundance of BLSC as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	3.8e-04		1.3e-03	9.1e-04
Rhode Island / Massachusetts OCS-A 0486	8.9e-05		3.1e-04	2.3e-04
Rhode Island / Massachusetts OCS-A 0487	3.8e-05		1.7e-04	1.2e-04
Massachusetts OCS-A 0500	1.1e-04		4.6e-04	3.3e-04
Massachusetts OCS-A 0501	9.2e-05		4.3e-04	3.3e-04
Massachusetts OCS-A 0502	1.1e-04		5.7e-04	5.0e-04
Massachusetts OCS-A 0503	5.9e-05		3.1e-04	3.2e-04
New York Proposed Commercial Lease - Unsolicited	2.7e-05		1.0e-04	1.0e-04
New York OCS-A 0512	1.1e-04		2.4e-04	3.2e-04
New Jersey OCS-A 0499	9.1e-04		5.1e-04	7.4e-04
New Jersey OCS-A 0498	5.0e-04		4.5e-04	6.9e-04
Delaware OCS-A 0482	1.1e-04		2.7e-04	4.7e-04
Maryland OCS-A 0489	6.2e-05		9.7e-05	1.5e-04
Maryland OCS-A 0490	4.7e-05		1.1e-04	1.5e-04
Virginia OCS-A 0483	5.2e-05		2.4e-04	2.6e-04
Virginia OCS-A 0497	2.6e-05		1.2e-05	2.2e-05
North Carolina OCS-A 0508	4.3e-05		2.4e-04	2.2e-04
North Carolina WEA - Wilmington West	6.4e-05		1.3e-04	4.0e-04
North Carolina WEA - Wilmington East	6.4e-05		2.9e-04	5.1e-04
South Carolina Call Area - Grand Strand	1.3e-03		1.6e-03	4.7e-03
South Carolina Call Area - Cape Romain	5.4e-04		4.5e-04	2.5e-03
South Carolina Call Area - Winyah	9.6e-06		7.1e-05	1.4e-04
South Carolina Call Area - Charleston	1.0e-05		7.4e-05	2.4e-04
All	4.7e-03		8.5e-03	1.4e-02

Predicted relative abundance of BOGU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	7.3e-05		6.2e-05	8.1e-05
Rhode Island / Massachusetts OCS-A 0486	9.2e-05		1.9e-04	8.9e-05
Rhode Island / Massachusetts OCS-A 0487	6.3e-05		1.3e-04	5.4e-05
Massachusetts OCS-A 0500	1.6e-04		3.5e-04	1.7e-04
Massachusetts OCS-A 0501	1.4e-04		3.2e-04	1.6e-04
Massachusetts OCS-A 0502	2.1e-04		4.4e-04	2.2e-04
Massachusetts OCS-A 0503	1.1e-04		2.4e-04	1.2e-04
New York Proposed Commercial Lease - Unsolicited	6.7e-05		8.2e-05	4.8e-05
New York OCS-A 0512	2.3e-04		1.7e-04	1.3e-04
New Jersey OCS-A 0499	1.2e-03		4.7e-04	3.2e-04
New Jersey OCS-A 0498	1.2e-03		4.4e-04	2.9e-04
Delaware OCS-A 0482	6.6e-04		2.8e-04	1.6e-04
Maryland OCS-A 0489	1.4e-04		1.0e-04	6.3e-05
Maryland OCS-A 0490	2.5e-04		1.3e-04	9.5e-05
Virginia OCS-A 0483	5.0e-04		2.6e-04	6.3e-04
Virginia OCS-A 0497	1.1e-05		7.4e-06	2.0e-05
North Carolina OCS-A 0508	7.6e-04		2.6e-04	5.7e-04
North Carolina WEA - Wilmington West	1.3e-03		2.4e-04	3.4e-04
North Carolina WEA - Wilmington East	7.4e-04		5.6e-04	1.5e-03
South Carolina Call Area - Grand Strand	1.0e-02		3.2e-03	2.6e-03
South Carolina Call Area - Cape Romain	1.7e-03		9.7e-04	5.2e-04
South Carolina Call Area - Winyah	6.9e-05		1.2e-04	5.7e-05
South Carolina Call Area - Charleston	9.8e-05		1.5e-04	1.3e-04
All	2.0e-02		9.2e-03	8.4e-03

Predicted relative abundance of BRPE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.4e-05	4.6e-08	8.7e-06	2.7e-08
Rhode Island / Massachusetts OCS-A 0486	8.0e-05	1.1e-07	2.7e-05	9.7e-08
Rhode Island / Massachusetts OCS-A 0487	5.4e-05	7.3e-08	1.9e-05	6.6e-08
Massachusetts OCS-A 0500	1.5e-04	1.9e-07	5.0e-05	1.8e-07
Massachusetts OCS-A 0501	1.3e-04	1.7e-07	4.3e-05	1.5e-07
Massachusetts OCS-A 0502	1.9e-04	2.4e-07	6.2e-05	2.2e-07
Massachusetts OCS-A 0503	1.1e-04	1.3e-07	3.4e-05	1.2e-07
New York Proposed Commercial Lease - Unsolicited	3.5e-05	5.8e-08	1.4e-05	5.4e-08
New York OCS-A 0512	6.9e-05	1.6e-07	3.1e-05	1.2e-07
New Jersey OCS-A 0499	1.7e-04	2.2e-06	7.3e-05	3.3e-07
New Jersey OCS-A 0498	1.5e-04	5.7e-07	6.4e-05	3.1e-07
Delaware OCS-A 0482	9.4e-05	2.9e-07	3.9e-05	2.3e-07
Maryland OCS-A 0489	3.3e-05	8.4e-08	1.3e-05	8.2e-08
Maryland OCS-A 0490	4.7e-05	8.6e-08	1.8e-05	1.2e-07
Virginia OCS-A 0483	1.3e-04	1.9e-07	4.3e-05	4.3e-07
Virginia OCS-A 0497	3.5e-06	7.9e-06	1.3e-06	1.6e-08
North Carolina OCS-A 0508	1.4e-04	1.7e-07	4.3e-05	4.3e-07
North Carolina WEA - Wilmington West	1.4e-04	1.9e-06	4.0e-05	1.3e-06
North Carolina WEA - Wilmington East	4.2e-04	2.4e-05	6.8e-05	2.6e-06
South Carolina Call Area - Grand Strand	2.1e-03	5.6e-05	7.9e-04	2.8e-05
South Carolina Call Area - Cape Romain	1.1e-03	1.5e-04	1.6e-04	6.8e-06
South Carolina Call Area - Winyah	8.7e-05	4.7e-08	1.5e-05	4.5e-07
South Carolina Call Area - Charleston	9.4e-05	5.1e-08	1.5e-05	1.4e-06
All	5.5e-03	2.5e-04	1.7e-03	4.3e-05

Predicted relative abundance of BRSP as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478		3.2e-06		
Rhode Island / Massachusetts OCS-A 0486		1.1e-05		
Rhode Island / Massachusetts OCS-A 0487		7.9e-06		
Massachusetts OCS-A 0500		2.2e-05		
Massachusetts OCS-A 0501		1.9e-05		
Massachusetts OCS-A 0502		2.9e-05		
Massachusetts OCS-A 0503		1.7e-05		
New York Proposed Commercial Lease - Unsolicited		5.2e-06		
New York OCS-A 0512		1.0e-05		
New Jersey OCS-A 0499		2.5e-05		
New Jersey OCS-A 0498		2.2e-05		
Delaware OCS-A 0482		1.3e-05		
Maryland OCS-A 0489		4.5e-06		
Maryland OCS-A 0490		6.5e-06		
Virginia OCS-A 0483		1.6e-05		
Virginia OCS-A 0497		3.8e-07		
North Carolina OCS-A 0508		1.9e-05		
North Carolina WEA - Wilmington West		6.3e-06		
North Carolina WEA - Wilmington East		1.7e-05		
South Carolina Call Area - Grand Strand		7.7e-05		
South Carolina Call Area - Cape Romain		1.9e-05		
South Carolina Call Area - Winyah		4.7e-06		
South Carolina Call Area - Charleston		4.4e-06		
All		3.6e-04		

Predicted relative abundance of BRTE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478		9.7e-06	2.2e-05	
Rhode Island / Massachusetts OCS-A 0486		3.3e-05	7.2e-05	
Rhode Island / Massachusetts OCS-A 0487		2.3e-05	5.0e-05	
Massachusetts OCS-A 0500		6.3e-05	1.4e-04	
Massachusetts OCS-A 0501		5.6e-05	1.2e-04	
Massachusetts OCS-A 0502		8.4e-05	1.8e-04	
Massachusetts OCS-A 0503		4.8e-05	1.0e-04	
New York Proposed Commercial Lease - Unsolicited		1.4e-05	3.1e-05	
New York OCS-A 0512		2.6e-05	5.9e-05	
New Jersey OCS-A 0499		6.2e-05	1.4e-04	
New Jersey OCS-A 0498		5.4e-05	1.2e-04	
Delaware OCS-A 0482		3.3e-05	7.1e-05	
Maryland OCS-A 0489		1.1e-05	2.4e-05	
Maryland OCS-A 0490		1.6e-05	3.5e-05	
Virginia OCS-A 0483		4.3e-05	8.8e-05	
Virginia OCS-A 0497		1.1e-06	2.1e-06	
North Carolina OCS-A 0508		5.1e-05	4.6e-04	
North Carolina WEA - Wilmington West		2.6e-05	4.3e-05	
North Carolina WEA - Wilmington East		4.4e-04	1.4e-04	
South Carolina Call Area - Grand Strand		4.6e-04	7.5e-04	
South Carolina Call Area - Cape Romain		1.9e-04	1.2e-04	
South Carolina Call Area - Winyah		2.4e-04	1.2e-04	
South Carolina Call Area - Charleston		9.6e-05	5.3e-05	
All		2.1e-03	2.9e-03	

Predicted relative abundance of COEI as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.2e-04	2.0e-05	5.7e-04	7.6e-04
Rhode Island / Massachusetts OCS-A 0486	1.8e-04	5.1e-05	3.3e-04	5.1e-04
Rhode Island / Massachusetts OCS-A 0487	1.2e-04	3.4e-05	2.2e-04	3.3e-04
Massachusetts OCS-A 0500	3.5e-04	9.0e-05	6.0e-04	9.9e-04
Massachusetts OCS-A 0501	3.2e-04	7.7e-05	5.3e-04	9.5e-04
Massachusetts OCS-A 0502	4.8e-04	1.1e-04	7.9e-04	1.4e-03
Massachusetts OCS-A 0503	2.8e-04	6.0e-05	4.7e-04	8.1e-04
New York Proposed Commercial Lease - Unsolicited	7.7e-05	2.0e-05	1.3e-04	1.1e-04
New York OCS-A 0512	1.5e-04	3.9e-05	2.7e-04	1.6e-04
New Jersey OCS-A 0499	3.8e-04	7.8e-05	5.9e-04	2.5e-04
New Jersey OCS-A 0498	3.4e-04	6.8e-05	5.1e-04	1.9e-04
Delaware OCS-A 0482	2.1e-04	4.1e-05	3.1e-04	9.6e-05
Maryland OCS-A 0489	7.2e-05	1.4e-05	1.1e-04	3.1e-05
Maryland OCS-A 0490	1.0e-04	1.9e-05	1.5e-04	4.5e-05
Virginia OCS-A 0483	2.7e-04	3.9e-05	3.7e-04	7.4e-05
Virginia OCS-A 0497	6.7e-06	9.8e-07	9.6e-06	1.7e-06
North Carolina OCS-A 0508	3.3e-04	3.6e-05	3.9e-04	8.2e-05
North Carolina WEA - Wilmington West	1.2e-04	1.5e-05	1.5e-04	1.1e-05
North Carolina WEA - Wilmington East	3.4e-04	3.7e-05	3.9e-04	3.1e-05
South Carolina Call Area - Grand Strand	1.5e-03	1.9e-04	1.7e-03	1.2e-04
South Carolina Call Area - Cape Romain	3.5e-04	6.4e-05	4.7e-04	2.5e-05
South Carolina Call Area - Winyah	8.6e-05	1.1e-05	9.7e-05	7.1e-06
South Carolina Call Area - Charleston	8.3e-05	1.2e-05	9.6e-05	5.9e-06
All	6.3e-03	1.1e-03	9.3e-03	7.0e-03

Predicted relative abundance of COLO as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.0e-03	1.7e-03	1.9e-03	1.5e-03
Rhode Island / Massachusetts OCS-A 0486	1.8e-03	5.1e-04	7.3e-04	1.9e-03
Rhode Island / Massachusetts OCS-A 0487	6.4e-04	2.3e-04	2.7e-04	5.4e-04
Massachusetts OCS-A 0500	1.2e-03	5.6e-04	5.5e-04	1.2e-03
Massachusetts OCS-A 0501	7.1e-04	5.0e-04	4.7e-04	9.5e-04
Massachusetts OCS-A 0502	7.2e-04	7.2e-04	5.0e-04	9.7e-04
Massachusetts OCS-A 0503	3.7e-04	3.8e-04	2.4e-04	5.4e-04
New York Proposed Commercial Lease - Unsolicited	4.8e-04	1.1e-04	3.0e-04	3.2e-04
New York OCS-A 0512	1.8e-03	7.1e-04	8.8e-04	1.0e-03
New Jersey OCS-A 0499	1.1e-02	3.6e-03	3.5e-03	4.8e-03
New Jersey OCS-A 0498	1.1e-02	1.6e-03	3.5e-03	4.1e-03
Delaware OCS-A 0482	3.1e-03	7.3e-04	1.1e-03	1.7e-03
Maryland OCS-A 0489	1.0e-03	2.8e-04	4.5e-04	7.2e-04
Maryland OCS-A 0490	1.4e-03	2.1e-04	4.8e-04	8.2e-04
Virginia OCS-A 0483	4.7e-03	2.0e-04	4.4e-04	1.8e-03
Virginia OCS-A 0497	1.4e-04	8.8e-06	1.4e-05	6.5e-05
North Carolina OCS-A 0508	2.3e-03	1.9e-04	3.2e-04	7.0e-04
North Carolina WEA - Wilmington West	8.2e-04	1.1e-04	2.1e-04	5.7e-04
North Carolina WEA - Wilmington East	8.6e-04	2.0e-04	3.0e-04	6.3e-04
South Carolina Call Area - Grand Strand	8.5e-03	1.9e-03	2.8e-03	7.4e-03
South Carolina Call Area - Cape Romain	1.6e-03	1.1e-03	8.5e-04	1.8e-03
South Carolina Call Area - Winyah	9.4e-05	5.1e-05	7.2e-05	1.6e-04
South Carolina Call Area - Charleston	1.6e-04	5.5e-05	9.1e-05	2.0e-04
All	5.5e-02	1.6e-02	2.0e-02	3.4e-02

Predicted relative abundance of COMU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	5.2e-05			3.8e-06
Rhode Island / Massachusetts OCS-A 0486	3.2e-04			3.1e-04
Rhode Island / Massachusetts OCS-A 0487	2.4e-04			1.2e-04
Massachusetts OCS-A 0500	7.1e-04			3.9e-04
Massachusetts OCS-A 0501	6.0e-04			3.9e-04
Massachusetts OCS-A 0502	9.6e-04			6.3e-04
Massachusetts OCS-A 0503	6.4e-04			4.0e-04
New York Proposed Commercial Lease - Unsolicited	8.6e-05			8.7e-05
New York OCS-A 0512	1.5e-04			1.7e-02
New Jersey OCS-A 0499	2.3e-04			4.8e-05
New Jersey OCS-A 0498	1.4e-04			6.5e-05
Delaware OCS-A 0482	7.0e-05			4.8e-06
Maryland OCS-A 0489	2.3e-05			1.6e-06
Maryland OCS-A 0490	3.3e-05			2.2e-06
Virginia OCS-A 0483	6.5e-05			5.6e-06
Virginia OCS-A 0497	1.5e-06			1.3e-07
North Carolina OCS-A 0508	7.3e-05			5.7e-06
North Carolina WEA - Wilmington West	2.0e-05			2.4e-06
North Carolina WEA - Wilmington East	5.6e-05			5.9e-06
South Carolina Call Area - Grand Strand	2.3e-04			2.8e-05
South Carolina Call Area - Cape Romain	5.5e-05			7.4e-06
South Carolina Call Area - Winyah	1.8e-05			1.3e-06
South Carolina Call Area - Charleston	1.5e-05			1.4e-06
All	4.8e-03			2.0e-02

Predicted relative abundance of COSH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.3e-05	8.7e-06	7.7e-06	
Rhode Island / Massachusetts OCS-A 0486	7.8e-05	9.5e-04	1.6e-04	
Rhode Island / Massachusetts OCS-A 0487	5.6e-05	1.0e-03	1.7e-04	
Massachusetts OCS-A 0500	1.6e-04	1.8e-03	5.1e-04	
Massachusetts OCS-A 0501	1.4e-04	1.5e-03	5.5e-04	
Massachusetts OCS-A 0502	2.2e-04	1.7e-03	8.4e-04	
Massachusetts OCS-A 0503	1.4e-04	1.1e-03	6.0e-04	
New York Proposed Commercial Lease - Unsolicited	3.2e-05	1.2e-04	2.9e-05	
New York OCS-A 0512	5.9e-05	1.7e-04	3.0e-05	
New Jersey OCS-A 0499	1.4e-04	2.3e-04	5.2e-05	
New Jersey OCS-A 0498	1.4e-04	1.8e-04	4.6e-05	
Delaware OCS-A 0482	8.0e-05	7.7e-05	2.6e-05	
Maryland OCS-A 0489	2.7e-05	2.6e-05	8.3e-06	
Maryland OCS-A 0490	4.0e-05	5.7e-05	1.4e-05	
Virginia OCS-A 0483	1.3e-04	1.3e-04	4.0e-05	
Virginia OCS-A 0497	2.7e-06	2.5e-06	7.9e-07	
North Carolina OCS-A 0508	5.0e-04	1.9e-04	1.2e-04	
North Carolina WEA - Wilmington West	5.0e-05	1.1e-04	2.0e-05	
North Carolina WEA - Wilmington East	2.3e-04	4.0e-04	2.5e-04	
South Carolina Call Area - Grand Strand	6.1e-04	9.5e-04	3.2e-04	
South Carolina Call Area - Cape Romain	1.6e-04	1.6e-04	8.4e-05	
South Carolina Call Area - Winyah	1.0e-04	9.4e-05	2.9e-04	
South Carolina Call Area - Charleston	5.1e-05	7.6e-05	2.2e-04	
All	3.2e-03	1.1e-02	4.4e-03	

Predicted relative abundance of COTE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.1e-03	3.7e-04	1.7e-04	
Rhode Island / Massachusetts OCS-A 0486	3.9e-04	5.4e-04	2.8e-04	
Rhode Island / Massachusetts OCS-A 0487	2.4e-04	2.6e-04	1.9e-04	
Massachusetts OCS-A 0500	5.9e-04	6.8e-04	5.2e-04	
Massachusetts OCS-A 0501	5.1e-04	5.7e-04	4.7e-04	
Massachusetts OCS-A 0502	6.7e-04	6.2e-04	7.4e-04	
Massachusetts OCS-A 0503	3.4e-04	3.3e-04	4.1e-04	
New York Proposed Commercial Lease - Unsolicited	3.3e-04	1.9e-04	1.2e-04	
New York OCS-A 0512	1.6e-03	1.3e-03	2.4e-04	
New Jersey OCS-A 0499	7.0e-03	1.5e-02	6.1e-04	
New Jersey OCS-A 0498	5.4e-03	7.4e-03	5.3e-04	
Delaware OCS-A 0482	1.8e-03	1.8e-03	2.9e-04	
Maryland OCS-A 0489	4.7e-04	5.0e-04	1.0e-04	
Maryland OCS-A 0490	6.7e-04	4.0e-04	1.3e-04	
Virginia OCS-A 0483	9.7e-04	2.9e-04	3.1e-04	
Virginia OCS-A 0497	3.2e-05	1.4e-05	8.5e-06	
North Carolina OCS-A 0508	5.7e-04	2.7e-04	3.9e-04	
North Carolina WEA - Wilmington West	2.9e-04	1.7e-04	2.0e-04	
North Carolina WEA - Wilmington East	3.5e-04	2.9e-04	4.9e-04	
South Carolina Call Area - Grand Strand	4.1e-03	2.3e-03	2.7e-03	
South Carolina Call Area - Cape Romain	1.2e-03	6.5e-04	7.0e-04	
South Carolina Call Area - Winyah	7.4e-05	6.2e-05	1.3e-04	
South Carolina Call Area - Charleston	1.4e-04	7.3e-05	1.5e-04	
All	2.9e-02	3.4e-02	9.8e-03	

Predicted relative abundance of DCCO as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.4e-04	3.6e-04	1.7e-04	8.0e-05
Rhode Island / Massachusetts OCS-A 0486	4.4e-04	5.1e-04	3.5e-04	2.1e-04
Rhode Island / Massachusetts OCS-A 0487	2.8e-04	2.9e-04	2.2e-04	1.4e-04
Massachusetts OCS-A 0500	6.6e-04	8.0e-04	5.5e-04	3.9e-04
Massachusetts OCS-A 0501	5.2e-04	7.0e-04	4.7e-04	3.5e-04
Massachusetts OCS-A 0502	7.3e-04	9.7e-04	6.4e-04	5.1e-04
Massachusetts OCS-A 0503	4.6e-04	5.5e-04	3.4e-04	2.9e-04
New York Proposed Commercial Lease - Unsolicited	2.0e-04	1.7e-04	1.6e-04	1.1e-04
New York OCS-A 0512	5.9e-04	3.5e-04	3.7e-04	2.0e-04
New Jersey OCS-A 0499	1.2e-03	6.8e-04	6.5e-04	5.2e-04
New Jersey OCS-A 0498	7.8e-04	5.5e-04	5.1e-04	4.9e-04
Delaware OCS-A 0482	4.5e-04	3.1e-04	2.9e-04	2.5e-04
Maryland OCS-A 0489	1.5e-04	1.0e-04	9.7e-05	8.6e-05
Maryland OCS-A 0490	1.7e-04	1.4e-04	1.2e-04	1.2e-04
Virginia OCS-A 0483	2.4e-04	2.6e-04	2.2e-04	3.0e-04
Virginia OCS-A 0497	2.3e-05	6.9e-06	1.1e-05	7.6e-06
North Carolina OCS-A 0508	1.7e-04	2.5e-04	2.0e-04	3.5e-04
North Carolina WEA - Wilmington West	1.3e-04	1.1e-04	7.5e-05	1.2e-04
North Carolina WEA - Wilmington East	2.6e-04	2.6e-04	1.7e-04	3.2e-04
South Carolina Call Area - Grand Strand	2.4e-03	1.4e-03	9.9e-04	1.6e-03
South Carolina Call Area - Cape Romain	9.6e-04	4.3e-04	2.5e-04	4.1e-04
South Carolina Call Area - Winyah	5.8e-05	6.7e-05	5.8e-05	9.5e-05
South Carolina Call Area - Charleston	8.2e-05	7.3e-05	4.9e-05	8.8e-05
All	1.1e-02	9.3e-03	7.0e-03	7.0e-03

Predicted relative abundance of DOVE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.1e-05	1.6e-05	5.8e-05	9.9e-06
Rhode Island / Massachusetts OCS-A 0486	4.8e-05	5.3e-05	1.8e-04	6.2e-05
Rhode Island / Massachusetts OCS-A 0487	3.9e-05	3.7e-05	1.2e-04	9.0e-05
Massachusetts OCS-A 0500	1.0e-04	1.1e-04	3.5e-04	2.1e-04
Massachusetts OCS-A 0501	8.8e-05	9.9e-05	3.2e-04	3.0e-04
Massachusetts OCS-A 0502	1.6e-04	2.3e-04	5.0e-04	4.6e-04
Massachusetts OCS-A 0503	1.2e-04	1.9e-04	2.9e-04	1.6e-04
New York Proposed Commercial Lease - Unsolicited	2.1e-05	2.2e-05	6.6e-05	4.6e-05
New York OCS-A 0512	3.8e-05	4.2e-05	1.4e-04	1.4e-04
New Jersey OCS-A 0499	6.0e-05	9.6e-05	3.4e-04	7.7e-05
New Jersey OCS-A 0498	4.3e-05	8.4e-05	3.0e-04	7.6e-05
Delaware OCS-A 0482	2.3e-05	5.0e-05	1.9e-04	5.4e-05
Maryland OCS-A 0489	7.6e-06	1.7e-05	6.3e-05	1.4e-05
Maryland OCS-A 0490	1.1e-05	2.5e-05	9.0e-05	3.4e-05
Virginia OCS-A 0483	2.5e-05	5.9e-05	2.2e-04	4.4e-05
Virginia OCS-A 0497	5.9e-07	1.4e-06	5.5e-06	8.6e-07
North Carolina OCS-A 0508	5.4e-05	6.4e-05	2.0e-04	1.2e-04
North Carolina WEA - Wilmington West	1.1e-05	2.7e-05	1.1e-04	1.8e-05
North Carolina WEA - Wilmington East	3.4e-05	6.9e-05	2.8e-04	5.6e-05
South Carolina Call Area - Grand Strand	1.3e-04	3.2e-04	1.3e-03	2.3e-04
South Carolina Call Area - Cape Romain	3.2e-05	8.2e-05	3.3e-04	6.1e-05
South Carolina Call Area - Winyah	1.4e-05	1.8e-05	6.2e-05	8.6e-06
South Carolina Call Area - Charleston	9.8e-06	1.8e-05	7.3e-05	1.2e-05
All	1.1e-03	1.7e-03	5.6e-03	2.3e-03

Predicted relative abundance of GBBG as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	3.1e-04	2.4e-04	3.3e-04	1.2e-04
Rhode Island / Massachusetts OCS-A 0486	1.0e-03	7.4e-04	6.4e-04	4.1e-04
Rhode Island / Massachusetts OCS-A 0487	6.8e-04	3.8e-04	3.6e-04	2.7e-04
Massachusetts OCS-A 0500	1.9e-03	8.8e-04	8.7e-04	6.8e-04
Massachusetts OCS-A 0501	1.7e-03	8.9e-04	7.7e-04	6.0e-04
Massachusetts OCS-A 0502	2.3e-03	8.5e-04	1.1e-03	8.6e-04
Massachusetts OCS-A 0503	1.1e-03	3.6e-04	6.3e-04	4.6e-04
New York Proposed Commercial Lease - Unsolicited	4.9e-04	1.1e-04	1.6e-04	1.8e-04
New York OCS-A 0512	8.6e-04	2.0e-04	3.0e-04	3.3e-04
New Jersey OCS-A 0499	1.5e-03	5.2e-04	6.4e-04	6.0e-04
New Jersey OCS-A 0498	1.1e-03	1.7e-04	5.5e-04	5.3e-04
Delaware OCS-A 0482	2.3e-04	5.4e-05	2.6e-04	2.6e-04
Maryland OCS-A 0489	6.5e-05	1.7e-05	9.4e-05	7.8e-05
Maryland OCS-A 0490	6.6e-05	1.5e-05	1.0e-04	1.2e-04
Virginia OCS-A 0483	8.8e-05	1.9e-05	1.5e-04	3.1e-04
Virginia OCS-A 0497	3.3e-06	6.4e-07	5.1e-06	8.2e-06
North Carolina OCS-A 0508	5.7e-05	1.7e-05	1.3e-04	3.0e-04
North Carolina WEA - Wilmington West	2.6e-05	7.7e-06	3.3e-05	6.2e-05
North Carolina WEA - Wilmington East	2.5e-05	1.6e-05	6.9e-05	1.7e-04
South Carolina Call Area - Grand Strand	2.3e-04	1.0e-04	3.6e-04	7.3e-04
South Carolina Call Area - Cape Romain	4.5e-05	3.1e-05	8.5e-05	1.8e-04
South Carolina Call Area - Winyah	3.9e-06	3.8e-06	1.0e-05	4.0e-05
South Carolina Call Area - Charleston	4.3e-06	4.3e-06	1.4e-05	4.0e-05
All	1.4e-02	5.6e-03	7.7e-03	7.3e-03

Predicted relative abundance of GRSH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.4e-05	1.7e-05	6.4e-06	1.7e-05
Rhode Island / Massachusetts OCS-A 0486	4.7e-05	2.5e-04	1.1e-04	4.8e-05
Rhode Island / Massachusetts OCS-A 0487	3.8e-05	2.2e-04	1.6e-04	3.3e-05
Massachusetts OCS-A 0500	1.2e-04	5.2e-04	5.2e-04	9.3e-05
Massachusetts OCS-A 0501	1.1e-04	4.4e-04	5.2e-04	8.6e-05
Massachusetts OCS-A 0502	2.2e-04	6.4e-04	1.1e-03	1.3e-04
Massachusetts OCS-A 0503	1.7e-04	4.0e-04	9.2e-04	7.8e-05
New York Proposed Commercial Lease - Unsolicited	1.1e-05	1.0e-04	3.1e-05	1.9e-05
New York OCS-A 0512	1.7e-05	9.9e-05	2.1e-05	3.6e-05
New Jersey OCS-A 0499	4.6e-05	1.1e-04	2.2e-05	9.1e-05
New Jersey OCS-A 0498	3.5e-05	1.0e-04	2.0e-05	8.1e-05
Delaware OCS-A 0482	2.1e-05	6.6e-05	1.3e-05	5.0e-05
Maryland OCS-A 0489	7.2e-06	2.0e-05	4.4e-06	1.8e-05
Maryland OCS-A 0490	1.0e-05	3.4e-05	7.3e-06	2.6e-05
Virginia OCS-A 0483	2.4e-05	7.4e-05	2.0e-05	7.2e-05
Virginia OCS-A 0497	5.9e-07	1.7e-06	3.9e-07	1.7e-06
North Carolina OCS-A 0508	2.4e-05	1.0e-04	2.2e-05	8.3e-05
North Carolina WEA - Wilmington West	1.5e-05	2.9e-05	6.3e-06	3.1e-05
North Carolina WEA - Wilmington East	4.4e-05	8.9e-05	2.1e-05	8.1e-05
South Carolina Call Area - Grand Strand	1.8e-04	3.5e-04	8.0e-05	3.7e-04
South Carolina Call Area - Cape Romain	4.2e-05	9.1e-05	1.9e-05	9.1e-05
South Carolina Call Area - Winyah	2.1e-05	3.3e-05	3.4e-06	2.5e-05
South Carolina Call Area - Charleston	9.7e-06	2.4e-05	5.5e-06	2.0e-05
All	1.2e-03	3.8e-03	3.7e-03	1.6e-03

Predicted relative abundance of GRSK as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478			1.7e-05	
Rhode Island / Massachusetts OCS-A 0486			5.2e-05	
Rhode Island / Massachusetts OCS-A 0487			3.9e-05	
Massachusetts OCS-A 0500			1.2e-04	
Massachusetts OCS-A 0501			1.0e-04	
Massachusetts OCS-A 0502			1.8e-04	
Massachusetts OCS-A 0503			1.3e-04	
New York Proposed Commercial Lease - Unsolicited			1.6e-05	
New York OCS-A 0512			3.0e-05	
New Jersey OCS-A 0499			6.4e-05	
New Jersey OCS-A 0498			5.6e-05	
Delaware OCS-A 0482			3.3e-05	
Maryland OCS-A 0489			1.1e-05	
Maryland OCS-A 0490			1.6e-05	
Virginia OCS-A 0483			4.0e-05	
Virginia OCS-A 0497			9.5e-07	
North Carolina OCS-A 0508			4.8e-05	
North Carolina WEA - Wilmington West			1.5e-05	
North Carolina WEA - Wilmington East			4.2e-05	
South Carolina Call Area - Grand Strand			1.8e-04	
South Carolina Call Area - Cape Romain			4.4e-05	
South Carolina Call Area - Winyah			1.6e-05	
South Carolina Call Area - Charleston			1.1e-05	
All			1.3e-03	

Predicted relative abundance of HERG as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.5e-04	1.4e-04	3.0e-04	1.6e-04
Rhode Island / Massachusetts OCS-A 0486	7.3e-04	2.4e-03	8.7e-04	4.0e-04
Rhode Island / Massachusetts OCS-A 0487	5.0e-04	1.7e-03	5.8e-04	2.4e-04
Massachusetts OCS-A 0500	1.2e-03	3.0e-03	1.3e-03	5.6e-04
Massachusetts OCS-A 0501	9.8e-04	2.2e-03	1.1e-03	5.1e-04
Massachusetts OCS-A 0502	1.3e-03	1.6e-03	1.5e-03	7.7e-04
Massachusetts OCS-A 0503	6.8e-04	6.1e-04	8.3e-04	3.9e-04
New York Proposed Commercial Lease - Unsolicited	5.1e-04	2.8e-04	3.0e-04	1.9e-04
New York OCS-A 0512	1.0e-03	5.6e-04	5.1e-04	4.6e-04
New Jersey OCS-A 0499	1.7e-03	3.9e-04	8.3e-04	6.3e-04
New Jersey OCS-A 0498	1.1e-03	2.0e-04	6.1e-04	4.9e-04
Delaware OCS-A 0482	4.6e-04	7.6e-05	3.1e-04	1.2e-04
Maryland OCS-A 0489	1.4e-04	2.9e-05	1.1e-04	3.8e-05
Maryland OCS-A 0490	1.9e-04	2.8e-05	1.4e-04	5.5e-05
Virginia OCS-A 0483	2.8e-04	2.7e-05	1.8e-04	1.2e-04
Virginia OCS-A 0497	7.1e-06	1.3e-06	5.2e-06	3.5e-06
North Carolina OCS-A 0508	2.7e-04	2.0e-05	2.0e-04	2.4e-04
North Carolina WEA - Wilmington West	2.1e-04	1.8e-05	5.5e-05	4.8e-05
North Carolina WEA - Wilmington East	1.7e-04	2.5e-05	1.1e-04	1.3e-04
South Carolina Call Area - Grand Strand	1.5e-03	2.8e-04	6.2e-04	5.1e-04
South Carolina Call Area - Cape Romain	1.8e-04	1.4e-04	1.6e-04	1.3e-04
South Carolina Call Area - Winyah	2.2e-05	5.3e-06	1.4e-05	1.1e-04
South Carolina Call Area - Charleston	2.5e-05	6.8e-06	2.3e-05	2.8e-05
All	1.3e-02	1.4e-02	1.1e-02	6.4e-03

Predicted relative abundance of HOCR as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478				4.0e-04
Rhode Island / Massachusetts OCS-A 0486				1.5e-04
Rhode Island / Massachusetts OCS-A 0487				1.1e-04
Massachusetts OCS-A 0500				3.0e-04
Massachusetts OCS-A 0501				2.6e-04
Massachusetts OCS-A 0502				4.1e-04
Massachusetts OCS-A 0503				2.4e-04
New York Proposed Commercial Lease - Unsolicited				6.9e-05
New York OCS-A 0512				1.3e-04
New Jersey OCS-A 0499				3.6e-04
New Jersey OCS-A 0498				3.3e-04
Delaware OCS-A 0482				2.1e-04
Maryland OCS-A 0489				7.2e-05
Maryland OCS-A 0490				1.0e-04
Virginia OCS-A 0483				3.0e-04
Virginia OCS-A 0497				8.4e-06
North Carolina OCS-A 0508				3.6e-04
North Carolina WEA - Wilmington West				2.3e-04
North Carolina WEA - Wilmington East				5.3e-04
South Carolina Call Area - Grand Strand				2.9e-03
South Carolina Call Area - Cape Romain				6.0e-04
South Carolina Call Area - Winyah				1.2e-04
South Carolina Call Area - Charleston				1.1e-04
All				8.3e-03

Predicted relative abundance of LAGU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.1e-05	3.3e-05	8.0e-05	1.6e-05
Rhode Island / Massachusetts OCS-A 0486	4.7e-05	9.9e-05	3.7e-04	3.0e-05
Rhode Island / Massachusetts OCS-A 0487	2.9e-05	5.6e-05	1.5e-04	1.9e-05
Massachusetts OCS-A 0500	7.1e-05	1.4e-04	3.0e-04	5.0e-05
Massachusetts OCS-A 0501	5.8e-05	1.1e-04	2.2e-04	4.4e-05
Massachusetts OCS-A 0502	7.9e-05	1.3e-04	2.2e-04	6.2e-05
Massachusetts OCS-A 0503	4.6e-05	6.0e-05	8.7e-05	3.4e-05
New York Proposed Commercial Lease - Unsolicited	6.9e-05	5.9e-05	9.9e-05	1.5e-05
New York OCS-A 0512	3.2e-04	1.3e-03	3.2e-04	3.6e-05
New Jersey OCS-A 0499	3.6e-03	1.9e-02	2.3e-03	1.2e-04
New Jersey OCS-A 0498	3.3e-03	1.1e-02	2.3e-03	1.2e-04
Delaware OCS-A 0482	1.1e-03	2.6e-03	9.6e-04	7.5e-05
Maryland OCS-A 0489	3.0e-04	6.2e-04	5.0e-04	2.8e-05
Maryland OCS-A 0490	3.7e-04	3.9e-04	4.4e-04	3.4e-05
Virginia OCS-A 0483	9.0e-04	1.9e-04	5.5e-04	1.0e-04
Virginia OCS-A 0497	4.9e-05	3.1e-05	4.9e-05	5.4e-06
North Carolina OCS-A 0508	6.7e-04	1.1e-04	2.6e-04	8.1e-05
North Carolina WEA - Wilmington West	3.5e-04	4.8e-04	4.7e-04	7.7e-05
North Carolina WEA - Wilmington East	1.9e-04	3.6e-04	6.1e-04	1.5e-04
South Carolina Call Area - Grand Strand	7.6e-03	6.8e-03	5.0e-03	9.7e-04
South Carolina Call Area - Cape Romain	3.2e-03	3.1e-03	2.3e-03	2.6e-04
South Carolina Call Area - Winyah	4.7e-05	3.6e-05	4.9e-05	2.4e-05
South Carolina Call Area - Charleston	8.5e-05	8.2e-05	1.4e-04	4.1e-05
All	2.3e-02	4.6e-02	1.8e-02	2.4e-03

Predicted relative abundance of LESP as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.2e-05	2.7e-06	3.2e-06	
Rhode Island / Massachusetts OCS-A 0486	4.2e-05	1.0e-05	1.7e-05	
Rhode Island / Massachusetts OCS-A 0487	3.5e-05	7.8e-06	2.0e-05	
Massachusetts OCS-A 0500	9.2e-05	2.3e-05	5.1e-05	
Massachusetts OCS-A 0501	7.7e-05	2.2e-05	4.5e-05	
Massachusetts OCS-A 0502	1.3e-04	3.4e-05	6.8e-05	
Massachusetts OCS-A 0503	8.8e-05	2.5e-05	4.2e-05	
New York Proposed Commercial Lease - Unsolicited	1.5e-05	4.6e-06	7.8e-06	
New York OCS-A 0512	2.6e-05	7.4e-06	1.5e-05	
New Jersey OCS-A 0499	5.4e-05	1.4e-05	1.4e-05	
New Jersey OCS-A 0498	4.3e-05	1.2e-05	1.2e-05	
Delaware OCS-A 0482	2.4e-05	7.1e-06	6.9e-06	
Maryland OCS-A 0489	8.0e-06	2.4e-06	2.3e-06	
Maryland OCS-A 0490	1.2e-05	3.6e-06	3.4e-06	
Virginia OCS-A 0483	2.5e-05	8.7e-06	8.2e-06	
Virginia OCS-A 0497	5.9e-07	1.9e-07	2.0e-07	
North Carolina OCS-A 0508	4.2e-05	1.0e-05	9.6e-06	
North Carolina WEA - Wilmington West	7.6e-06	2.6e-06	3.5e-06	
North Carolina WEA - Wilmington East	2.1e-05	7.3e-06	9.3e-06	
South Carolina Call Area - Grand Strand	9.0e-05	3.2e-05	4.2e-05	
South Carolina Call Area - Cape Romain	2.1e-05	7.4e-06	1.1e-05	
South Carolina Call Area - Winyah	8.8e-06	2.3e-06	2.7e-06	
South Carolina Call Area - Charleston	5.2e-06	1.9e-06	2.5e-06	
All	8.8e-04	2.5e-04	4.0e-04	

Predicted relative abundance of LETE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478		1.8e-04	2.5e-07	
Rhode Island / Massachusetts OCS-A 0486		1.9e-04	4.5e-07	
Rhode Island / Massachusetts OCS-A 0487		1.3e-04	3.0e-07	
Massachusetts OCS-A 0500		3.5e-04	8.1e-07	
Massachusetts OCS-A 0501		3.1e-04	7.0e-07	
Massachusetts OCS-A 0502		4.4e-04	9.8e-07	
Massachusetts OCS-A 0503		2.4e-04	5.3e-07	
New York Proposed Commercial Lease - Unsolicited		8.3e-05	1.9e-07	
New York OCS-A 0512		1.7e-04	3.8e-07	
New Jersey OCS-A 0499		4.0e-04	8.5e-07	
New Jersey OCS-A 0498		3.2e-04	7.5e-07	
Delaware OCS-A 0482		1.9e-04	4.6e-07	
Maryland OCS-A 0489		6.4e-05	1.6e-07	
Maryland OCS-A 0490		8.8e-05	2.2e-07	
Virginia OCS-A 0483		2.0e-04	5.6e-07	
Virginia OCS-A 0497		5.9e-06	1.6e-08	
North Carolina OCS-A 0508		2.0e-04	5.6e-07	
North Carolina WEA - Wilmington West		2.3e-04	1.6e-06	
North Carolina WEA - Wilmington East		5.3e-04	4.8e-06	
South Carolina Call Area - Grand Strand		3.6e-03	1.2e-03	
South Carolina Call Area - Cape Romain		1.3e-03	8.9e-04	
South Carolina Call Area - Winyah		1.2e-04	2.7e-05	
South Carolina Call Area - Charleston		2.1e-04	1.8e-04	
All		9.6e-03	2.3e-03	

Predicted relative abundance of LTDU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	9.2e-03		8.6e-04	5.5e-03
Rhode Island / Massachusetts OCS-A 0486	1.1e-04		2.7e-04	2.8e-04
Rhode Island / Massachusetts OCS-A 0487	7.4e-05		1.7e-04	1.3e-04
Massachusetts OCS-A 0500	2.2e-04		4.1e-04	4.5e-04
Massachusetts OCS-A 0501	2.2e-04		3.6e-04	5.8e-04
Massachusetts OCS-A 0502	3.9e-04		4.8e-04	9.1e-04
Massachusetts OCS-A 0503	4.7e-03		2.6e-04	1.5e-03
New York Proposed Commercial Lease - Unsolicited	4.1e-05		1.0e-04	7.0e-05
New York OCS-A 0512	1.1e-04		2.0e-04	2.8e-04
New Jersey OCS-A 0499	1.8e-04		4.4e-04	4.8e-04
New Jersey OCS-A 0498	1.3e-04		3.8e-04	3.2e-04
Delaware OCS-A 0482	7.7e-05		2.4e-04	1.7e-04
Maryland OCS-A 0489	2.3e-05		8.3e-05	4.9e-05
Maryland OCS-A 0490	3.3e-05		1.1e-04	6.2e-05
Virginia OCS-A 0483	8.0e-05		2.5e-04	1.3e-04
Virginia OCS-A 0497	3.1e-06		6.4e-06	3.3e-06
North Carolina OCS-A 0508	8.0e-05		2.6e-04	1.4e-04
North Carolina WEA - Wilmington West	3.1e-05		1.3e-04	4.2e-05
North Carolina WEA - Wilmington East	8.0e-05		3.0e-04	1.3e-04
South Carolina Call Area - Grand Strand	4.6e-04		1.6e-03	5.0e-04
South Carolina Call Area - Cape Romain	1.1e-04		4.1e-04	1.1e-04
South Carolina Call Area - Winyah	2.0e-05		7.5e-05	3.0e-05
South Carolina Call Area - Charleston	2.0e-05		8.4e-05	2.8e-05
All	1.6e-02		7.4e-03	1.2e-02

Predicted relative abundance of MASH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.1e-06	5.9e-05	1.5e-05	
Rhode Island / Massachusetts OCS-A 0486	4.3e-06	3.2e-04	6.3e-05	
Rhode Island / Massachusetts OCS-A 0487	3.2e-06	3.0e-04	6.0e-05	
Massachusetts OCS-A 0500	9.2e-06	7.8e-04	1.5e-04	
Massachusetts OCS-A 0501	8.3e-06	6.9e-04	1.2e-04	
Massachusetts OCS-A 0502	1.4e-05	1.1e-03	2.0e-04	
Massachusetts OCS-A 0503	9.5e-06	6.4e-04	1.3e-04	
New York Proposed Commercial Lease - Unsolicited	2.0e-06	1.2e-04	6.9e-05	
New York OCS-A 0512	3.4e-06	1.6e-04	9.5e-05	
New Jersey OCS-A 0499	8.3e-06	2.6e-04	9.2e-05	
New Jersey OCS-A 0498	9.0e-06	2.3e-04	7.8e-05	
Delaware OCS-A 0482	5.4e-06	1.3e-04	4.8e-05	
Maryland OCS-A 0489	1.6e-06	4.3e-05	1.4e-05	
Maryland OCS-A 0490	2.5e-06	6.6e-05	2.4e-05	
Virginia OCS-A 0483	9.9e-06	1.4e-04	6.2e-05	
Virginia OCS-A 0497	2.0e-07	3.3e-06	1.1e-06	
North Carolina OCS-A 0508	3.7e-05	1.6e-04	9.9e-05	
North Carolina WEA - Wilmington West	2.1e-06	5.1e-05	1.3e-05	
North Carolina WEA - Wilmington East	7.4e-06	1.4e-04	4.1e-05	
South Carolina Call Area - Grand Strand	2.3e-05	6.2e-04	1.8e-04	
South Carolina Call Area - Cape Romain	5.4e-06	1.8e-04	4.2e-05	
South Carolina Call Area - Winyah	2.2e-06	3.9e-05	3.4e-05	
South Carolina Call Area - Charleston	1.6e-06	3.6e-05	1.2e-05	
All	1.7e-04	6.3e-03	1.6e-03	

Predicted relative abundance of NOFU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	7.6e-06	5.2e-05	7.3e-11	4.8e-07
Rhode Island / Massachusetts OCS-A 0486	5.3e-05	9.9e-05	2.9e-07	7.2e-06
Rhode Island / Massachusetts OCS-A 0487	8.1e-05	6.6e-05	9.1e-06	2.0e-05
Massachusetts OCS-A 0500	2.7e-04	2.0e-04	1.5e-04	3.2e-04
Massachusetts OCS-A 0501	2.0e-04	2.4e-04	1.4e-04	4.8e-04
Massachusetts OCS-A 0502	4.2e-04	3.4e-04	2.4e-04	6.2e-04
Massachusetts OCS-A 0503	2.4e-04	2.4e-04	2.6e-04	3.7e-04
New York Proposed Commercial Lease - Unsolicited	1.5e-05	2.1e-05	1.6e-08	3.1e-06
New York OCS-A 0512	1.9e-05	4.1e-05	3.5e-09	6.6e-06
New Jersey OCS-A 0499	2.9e-05	9.6e-05	5.8e-10	3.1e-06
New Jersey OCS-A 0498	2.4e-05	9.5e-05	4.2e-10	2.6e-06
Delaware OCS-A 0482	1.3e-05	6.9e-05	2.4e-10	1.5e-06
Maryland OCS-A 0489	4.4e-06	2.5e-05	8.1e-11	5.4e-07
Maryland OCS-A 0490	6.5e-06	3.3e-05	1.3e-10	9.4e-07
Virginia OCS-A 0483	1.5e-05	8.6e-05	3.0e-10	2.0e-06
Virginia OCS-A 0497	3.5e-07	2.1e-06	6.6e-12	4.8e-08
North Carolina OCS-A 0508	2.2e-05	5.4e-05	6.2e-10	5.3e-06
North Carolina WEA - Wilmington West	3.6e-06	4.9e-05	1.1e-10	7.5e-07
North Carolina WEA - Wilmington East	1.2e-05	1.1e-04	3.2e-10	2.6e-06
South Carolina Call Area - Grand Strand	4.1e-05	6.1e-04	1.4e-09	9.7e-06
South Carolina Call Area - Cape Romain	9.2e-06	1.6e-04	3.7e-10	2.3e-06
South Carolina Call Area - Winyah	3.4e-06	1.9e-05	1.2e-10	8.1e-07
South Carolina Call Area - Charleston	2.8e-06	3.0e-05	9.0e-11	6.9e-07
All	1.5e-03	2.7e-03	8.0e-04	1.9e-03

Predicted relative abundance of NOGA as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.5e-04	4.2e-05	1.3e-04	2.3e-05
Rhode Island / Massachusetts OCS-A 0486	7.8e-04	1.4e-04	1.0e-03	6.5e-04
Rhode Island / Massachusetts OCS-A 0487	5.3e-04	7.7e-05	8.0e-04	4.3e-04
Massachusetts OCS-A 0500	1.5e-03	2.0e-04	2.0e-03	9.6e-04
Massachusetts OCS-A 0501	1.3e-03	1.7e-04	1.9e-03	8.3e-04
Massachusetts OCS-A 0502	2.0e-03	2.2e-04	2.9e-03	1.1e-03
Massachusetts OCS-A 0503	1.1e-03	1.2e-04	1.7e-03	6.8e-04
New York Proposed Commercial Lease - Unsolicited	3.5e-04	5.5e-05	2.9e-04	3.6e-04
New York OCS-A 0512	7.3e-04	3.5e-04	4.7e-04	9.4e-04
New Jersey OCS-A 0499	2.2e-03	1.9e-03	1.2e-03	1.9e-03
New Jersey OCS-A 0498	1.9e-03	6.6e-04	1.5e-03	1.7e-03
Delaware OCS-A 0482	6.9e-04	2.1e-04	3.5e-04	9.7e-04
Maryland OCS-A 0489	2.3e-04	5.1e-05	1.0e-04	4.6e-04
Maryland OCS-A 0490	3.0e-04	5.1e-05	9.1e-05	7.6e-04
Virginia OCS-A 0483	1.6e-03	5.8e-05	6.7e-05	3.1e-03
Virginia OCS-A 0497	6.2e-05	2.3e-06	3.9e-06	1.9e-04
North Carolina OCS-A 0508	1.2e-03	4.7e-05	6.7e-05	1.2e-03
North Carolina WEA - Wilmington West	6.0e-04	4.4e-05	2.1e-05	8.7e-04
North Carolina WEA - Wilmington East	3.3e-04	7.1e-05	4.0e-05	2.0e-03
South Carolina Call Area - Grand Strand	5.9e-03	1.2e-03	2.2e-04	6.3e-03
South Carolina Call Area - Cape Romain	2.0e-03	1.6e-03	6.3e-05	2.1e-03
South Carolina Call Area - Winyah	1.6e-05	2.1e-05	7.9e-06	2.2e-05
South Carolina Call Area - Charleston	1.3e-05	2.9e-05	9.1e-06	1.4e-04
All	2.6e-02	7.3e-03	1.5e-02	2.8e-02

Predicted relative abundance of PAJA as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	5.6e-05	7.7e-05	7.0e-05	
Rhode Island / Massachusetts OCS-A 0486	1.8e-04	3.2e-04	2.3e-04	
Rhode Island / Massachusetts OCS-A 0487	1.3e-04	2.3e-04	1.6e-04	
Massachusetts OCS-A 0500	3.4e-04	6.7e-04	4.4e-04	
Massachusetts OCS-A 0501	3.0e-04	6.4e-04	3.8e-04	
Massachusetts OCS-A 0502	4.6e-04	9.9e-04	5.8e-04	
Massachusetts OCS-A 0503	2.6e-04	6.5e-04	3.4e-04	
New York Proposed Commercial Lease - Unsolicited	8.8e-05	1.1e-04	1.1e-04	
New York OCS-A 0512	1.8e-04	1.9e-04	2.1e-04	
New Jersey OCS-A 0499	4.7e-04	3.5e-04	4.6e-04	
New Jersey OCS-A 0498	4.2e-04	2.9e-04	4.0e-04	
Delaware OCS-A 0482	2.7e-04	1.7e-04	2.1e-04	
Maryland OCS-A 0489	9.1e-05	5.6e-05	7.3e-05	
Maryland OCS-A 0490	1.3e-04	8.1e-05	1.0e-04	
Virginia OCS-A 0483	3.1e-04	1.8e-04	2.1e-04	
Virginia OCS-A 0497	7.9e-06	4.5e-06	5.2e-06	
North Carolina OCS-A 0508	3.6e-04	2.0e-04	2.2e-04	
North Carolina WEA - Wilmington West	1.4e-04	7.2e-05	7.2e-05	
North Carolina WEA - Wilmington East	3.6e-04	1.9e-04	1.9e-04	
South Carolina Call Area - Grand Strand	1.8e-03	8.6e-04	8.8e-04	
South Carolina Call Area - Cape Romain	5.0e-04	2.1e-04	2.1e-04	
South Carolina Call Area - Winyah	1.1e-04	4.8e-05	5.1e-05	
South Carolina Call Area - Charleston	1.0e-04	4.7e-05	4.7e-05	
All	7.1e-03	6.6e-03	5.7e-03	

Predicted relative abundance of POJA as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	8.9e-06	4.0e-05	1.3e-05	
Rhode Island / Massachusetts OCS-A 0486	3.1e-05	1.8e-04	1.1e-04	
Rhode Island / Massachusetts OCS-A 0487	2.3e-05	1.4e-04	1.0e-04	
Massachusetts OCS-A 0500	6.5e-05	4.1e-04	2.9e-04	
Massachusetts OCS-A 0501	5.6e-05	3.7e-04	2.8e-04	
Massachusetts OCS-A 0502	8.9e-05	5.8e-04	4.8e-04	
Massachusetts OCS-A 0503	5.5e-05	3.6e-04	3.9e-04	
New York Proposed Commercial Lease - Unsolicited	1.4e-05	6.3e-05	3.1e-05	
New York OCS-A 0512	2.6e-05	9.8e-05	4.8e-05	
New Jersey OCS-A 0499	5.9e-05	1.9e-04	6.8e-05	
New Jersey OCS-A 0498	5.2e-05	1.6e-04	5.8e-05	
Delaware OCS-A 0482	3.2e-05	9.7e-05	3.4e-05	
Maryland OCS-A 0489	1.1e-05	3.2e-05	1.1e-05	
Maryland OCS-A 0490	1.6e-05	4.7e-05	1.6e-05	
Virginia OCS-A 0483	4.2e-05	1.2e-04	3.9e-05	
Virginia OCS-A 0497	1.0e-06	2.8e-06	8.8e-07	
North Carolina OCS-A 0508	5.8e-05	1.4e-04	4.9e-05	
North Carolina WEA - Wilmington West	2.0e-05	5.6e-05	1.5e-05	
North Carolina WEA - Wilmington East	7.6e-05	1.7e-04	4.4e-05	
South Carolina Call Area - Grand Strand	2.5e-04	6.8e-04	1.9e-04	
South Carolina Call Area - Cape Romain	6.3e-05	1.7e-04	4.8e-05	
South Carolina Call Area - Winyah	3.1e-05	4.9e-05	3.4e-05	
South Carolina Call Area - Charleston	2.1e-05	4.3e-05	1.4e-05	
All	1.1e-03	4.2e-03	2.4e-03	

Predicted relative abundance of RAZO as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.4e-03	4.5e-05	3.5e-04	1.6e-03
Rhode Island / Massachusetts OCS-A 0486	1.4e-03	1.4e-04	2.6e-04	7.0e-04
Rhode Island / Massachusetts OCS-A 0487	1.6e-03	9.1e-05	1.7e-04	4.1e-04
Massachusetts OCS-A 0500	8.3e-03	2.4e-04	4.4e-04	3.2e-03
Massachusetts OCS-A 0501	1.0e-02	2.1e-04	3.9e-04	5.2e-03
Massachusetts OCS-A 0502	1.8e-02	3.0e-04	5.3e-04	5.3e-03
Massachusetts OCS-A 0503	9.8e-03	1.6e-04	2.9e-04	3.4e-03
New York Proposed Commercial Lease - Unsolicited	2.3e-04	5.4e-05	9.2e-05	1.0e-04
New York OCS-A 0512	4.5e-04	1.1e-04	1.8e-04	2.5e-04
New Jersey OCS-A 0499	1.5e-03	2.2e-04	3.8e-04	8.2e-04
New Jersey OCS-A 0498	8.3e-04	1.9e-04	3.4e-04	6.5e-04
Delaware OCS-A 0482	2.1e-04	1.1e-04	2.0e-04	2.5e-04
Maryland OCS-A 0489	5.8e-05	3.7e-05	6.9e-05	8.8e-05
Maryland OCS-A 0490	8.7e-05	5.2e-05	9.9e-05	1.3e-04
Virginia OCS-A 0483	1.8e-04	1.1e-04	2.4e-04	2.9e-04
Virginia OCS-A 0497	4.3e-06	2.9e-06	5.8e-06	7.9e-06
North Carolina OCS-A 0508	1.9e-04	1.1e-04	2.5e-04	2.7e-04
North Carolina WEA - Wilmington West	8.6e-05	5.7e-05	1.2e-04	3.8e-05
North Carolina WEA - Wilmington East	1.6e-04	1.4e-04	2.8e-04	1.3e-04
South Carolina Call Area - Grand Strand	9.5e-04	7.3e-04	1.4e-03	4.9e-04
South Carolina Call Area - Cape Romain	2.1e-04	1.9e-04	3.8e-04	1.0e-04
South Carolina Call Area - Winyah	4.2e-05	3.8e-05	7.3e-05	3.3e-05
South Carolina Call Area - Charleston	4.0e-05	4.3e-05	7.9e-05	3.1e-05
All	5.6e-02	3.4e-03	6.6e-03	2.4e-02

Predicted relative abundance of RBGU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	7.5e-05	2.1e-04	1.7e-04	8.8e-05
Rhode Island / Massachusetts OCS-A 0486	2.7e-04	2.8e-04	5.4e-04	2.6e-04
Rhode Island / Massachusetts OCS-A 0487	1.7e-04	1.9e-04	3.3e-04	1.7e-04
Massachusetts OCS-A 0500	4.1e-04	5.3e-04	7.6e-04	4.6e-04
Massachusetts OCS-A 0501	3.2e-04	4.6e-04	6.1e-04	4.1e-04
Massachusetts OCS-A 0502	4.1e-04	6.8e-04	8.5e-04	6.0e-04
Massachusetts OCS-A 0503	2.1e-04	3.8e-04	4.3e-04	3.4e-04
New York Proposed Commercial Lease - Unsolicited	1.7e-04	1.3e-04	3.5e-04	1.2e-04
New York OCS-A 0512	4.3e-04	2.8e-04	8.5e-04	2.8e-04
New Jersey OCS-A 0499	8.4e-04	7.4e-04	1.6e-03	6.8e-04
New Jersey OCS-A 0498	6.5e-04	5.9e-04	1.2e-03	5.9e-04
Delaware OCS-A 0482	3.7e-04	3.5e-04	6.8e-04	3.5e-04
Maryland OCS-A 0489	1.2e-04	1.3e-04	2.3e-04	1.2e-04
Maryland OCS-A 0490	1.5e-04	1.6e-04	2.7e-04	1.6e-04
Virginia OCS-A 0483	2.5e-04	3.4e-04	4.3e-04	3.7e-04
Virginia OCS-A 0497	1.1e-05	9.6e-06	1.3e-05	1.2e-05
North Carolina OCS-A 0508	1.9e-04	3.5e-04	3.9e-04	3.2e-04
North Carolina WEA - Wilmington West	3.1e-04	1.5e-04	1.8e-04	3.3e-04
North Carolina WEA - Wilmington East	4.0e-04	3.6e-04	2.9e-04	5.3e-04
South Carolina Call Area - Grand Strand	6.6e-03	1.9e-03	2.3e-03	4.4e-03
South Carolina Call Area - Cape Romain	6.3e-03	5.5e-04	5.6e-04	1.2e-03
South Carolina Call Area - Winyah	9.2e-05	9.4e-05	7.0e-05	1.2e-04
South Carolina Call Area - Charleston	1.4e-04	9.5e-05	8.0e-05	1.3e-04
All	1.9e-02	9.0e-03	1.3e-02	1.2e-02

Predicted relative abundance of RBME as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	3.4e-04			2.3e-04
Rhode Island / Massachusetts OCS-A 0486	2.7e-04			4.5e-04
Rhode Island / Massachusetts OCS-A 0487	1.8e-04			2.9e-04
Massachusetts OCS-A 0500	4.6e-04			8.4e-04
Massachusetts OCS-A 0501	4.3e-04			7.6e-04
Massachusetts OCS-A 0502	7.5e-04			1.1e-03
Massachusetts OCS-A 0503	6.1e-04			6.3e-04
New York Proposed Commercial Lease - Unsolicited	1.2e-04			1.8e-04
New York OCS-A 0512	3.1e-04			3.9e-04
New Jersey OCS-A 0499	7.5e-04			8.1e-04
New Jersey OCS-A 0498	5.7e-04			7.1e-04
Delaware OCS-A 0482	3.8e-04			4.1e-04
Maryland OCS-A 0489	1.3e-04			1.4e-04
Maryland OCS-A 0490	1.3e-04			1.8e-04
Virginia OCS-A 0483	2.3e-04			3.8e-04
Virginia OCS-A 0497	9.9e-06			1.0e-05
North Carolina OCS-A 0508	2.0e-04			3.7e-04
North Carolina WEA - Wilmington West	1.1e-04			1.8e-04
North Carolina WEA - Wilmington East	2.1e-04			4.2e-04
South Carolina Call Area - Grand Strand	1.6e-03			2.3e-03
South Carolina Call Area - Cape Romain	7.0e-04			5.9e-04
South Carolina Call Area - Winyah	4.8e-05			1.1e-04
South Carolina Call Area - Charleston	5.4e-05			1.2e-04
All	8.6e-03			1.2e-02

Predicted relative abundance of REPH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	6.4e-05	6.7e-05	2.8e-05	
Rhode Island / Massachusetts OCS-A 0486	2.2e-04	2.1e-04	9.2e-05	
Rhode Island / Massachusetts OCS-A 0487	1.6e-04	1.5e-04	6.7e-05	
Massachusetts OCS-A 0500	4.6e-04	4.2e-04	2.1e-04	
Massachusetts OCS-A 0501	4.3e-04	3.8e-04	1.9e-04	
Massachusetts OCS-A 0502	6.6e-04	5.6e-04	3.6e-04	
Massachusetts OCS-A 0503	3.6e-04	3.2e-04	2.5e-04	
New York Proposed Commercial Lease - Unsolicited	7.8e-05	8.6e-05	3.2e-05	
New York OCS-A 0512	1.4e-04	1.6e-04	5.7e-05	
New Jersey OCS-A 0499	3.1e-04	3.8e-04	1.2e-04	
New Jersey OCS-A 0498	2.7e-04	3.4e-04	1.1e-04	
Delaware OCS-A 0482	1.6e-04	2.0e-04	6.3e-05	
Maryland OCS-A 0489	4.8e-05	6.9e-05	2.1e-05	
Maryland OCS-A 0490	7.7e-05	9.9e-05	3.2e-05	
Virginia OCS-A 0483	2.1e-04	2.5e-04	8.4e-05	
Virginia OCS-A 0497	5.1e-06	6.0e-06	1.9e-06	
North Carolina OCS-A 0508	2.1e-04	2.7e-04	1.1e-04	
North Carolina WEA - Wilmington West	9.2e-05	1.2e-04	3.2e-05	
North Carolina WEA - Wilmington East	3.2e-04	3.0e-04	1.0e-04	
South Carolina Call Area - Grand Strand	1.1e-03	1.4e-03	3.8e-04	
South Carolina Call Area - Cape Romain	2.7e-04	3.4e-04	9.4e-05	
South Carolina Call Area - Winyah	1.0e-04	8.0e-05	3.4e-05	
South Carolina Call Area - Charleston	7.9e-05	8.2e-05	2.5e-05	
All	5.8e-03	6.3e-03	2.5e-03	

Predicted relative abundance of RNPH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	5.6e-05	5.8e-05	3.3e-05	
Rhode Island / Massachusetts OCS-A 0486	1.9e-04	1.9e-04	1.3e-04	
Rhode Island / Massachusetts OCS-A 0487	1.3e-04	1.3e-04	9.4e-05	
Massachusetts OCS-A 0500	3.8e-04	3.4e-04	2.6e-04	
Massachusetts OCS-A 0501	3.3e-04	3.1e-04	2.2e-04	
Massachusetts OCS-A 0502	5.3e-04	4.4e-04	3.5e-04	
Massachusetts OCS-A 0503	3.7e-04	2.5e-04	2.2e-04	
New York Proposed Commercial Lease - Unsolicited	8.4e-05	8.0e-05	5.2e-05	
New York OCS-A 0512	1.6e-04	1.6e-04	9.8e-05	
New Jersey OCS-A 0499	3.4e-04	3.3e-04	2.1e-04	
New Jersey OCS-A 0498	2.9e-04	2.9e-04	1.9e-04	
Delaware OCS-A 0482	1.7e-04	1.7e-04	1.1e-04	
Maryland OCS-A 0489	5.9e-05	5.7e-05	3.7e-05	
Maryland OCS-A 0490	8.6e-05	8.0e-05	6.1e-05	
Virginia OCS-A 0483	2.0e-04	1.8e-04	1.8e-04	
Virginia OCS-A 0497	4.6e-06	4.5e-06	4.1e-06	
North Carolina OCS-A 0508	2.7e-04	1.8e-04	2.7e-04	
North Carolina WEA - Wilmington West	8.4e-05	7.9e-05	5.5e-05	
North Carolina WEA - Wilmington East	2.2e-04	1.9e-04	1.7e-04	
South Carolina Call Area - Grand Strand	1.0e-03	9.9e-04	7.1e-04	
South Carolina Call Area - Cape Romain	2.5e-04	2.5e-04	1.6e-04	
South Carolina Call Area - Winyah	7.1e-05	5.3e-05	7.1e-05	
South Carolina Call Area - Charleston	5.8e-05	5.7e-05	4.8e-05	
All	5.4e-03	4.9e-03	3.7e-03	

Predicted relative abundance of ROST as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	3.5e-04	5.0e-03	1.1e-04	
Rhode Island / Massachusetts OCS-A 0486	3.1e-04	3.4e-06	2.9e-04	
Rhode Island / Massachusetts OCS-A 0487	2.2e-04	2.0e-06	2.0e-04	
Massachusetts OCS-A 0500	5.9e-04	5.9e-06	5.6e-04	
Massachusetts OCS-A 0501	5.2e-04	5.0e-06	5.0e-04	
Massachusetts OCS-A 0502	7.6e-04	7.1e-06	7.5e-04	
Massachusetts OCS-A 0503	4.4e-04	4.2e-06	4.3e-04	
New York Proposed Commercial Lease - Unsolicited	1.4e-04	9.8e-07	1.2e-04	
New York OCS-A 0512	2.8e-04	1.9e-06	2.3e-04	
New Jersey OCS-A 0499	7.1e-04	4.4e-06	5.3e-04	
New Jersey OCS-A 0498	6.0e-04	3.6e-06	4.6e-04	
Delaware OCS-A 0482	3.8e-04	2.0e-06	2.8e-04	
Maryland OCS-A 0489	1.3e-04	7.9e-07	9.4e-05	
Maryland OCS-A 0490	1.8e-04	9.5e-07	1.3e-04	
Virginia OCS-A 0483	3.9e-04	2.0e-06	3.2e-04	
Virginia OCS-A 0497	1.0e-05	7.4e-08	8.0e-06	
North Carolina OCS-A 0508	3.9e-04	2.0e-06	3.5e-04	
North Carolina WEA - Wilmington West	1.9e-04	1.1e-06	1.5e-04	
North Carolina WEA - Wilmington East	4.0e-04	2.3e-05	3.8e-04	
South Carolina Call Area - Grand Strand	2.3e-03	1.8e-05	1.8e-03	
South Carolina Call Area - Cape Romain	7.3e-04	5.6e-05	4.6e-04	
South Carolina Call Area - Winyah	1.1e-04	5.3e-07	9.9e-05	
South Carolina Call Area - Charleston	1.1e-04	5.7e-07	1.0e-04	
All	1.0e-02	5.1e-03	8.4e-03	

Predicted relative abundance of ROYT as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.5e-07	2.0e-06	1.2e-05	
Rhode Island / Massachusetts OCS-A 0486	7.5e-07	7.2e-06	2.6e-05	
Rhode Island / Massachusetts OCS-A 0487	5.0e-07	5.2e-06	1.8e-05	
Massachusetts OCS-A 0500	1.3e-06	1.4e-05	4.7e-05	
Massachusetts OCS-A 0501	1.1e-06	1.2e-05	4.0e-05	
Massachusetts OCS-A 0502	1.6e-06	1.7e-05	5.7e-05	
Massachusetts OCS-A 0503	8.4e-07	9.6e-06	3.1e-05	
New York Proposed Commercial Lease - Unsolicited	4.4e-07	3.7e-06	1.7e-05	
New York OCS-A 0512	9.6e-07	8.6e-06	4.2e-05	
New Jersey OCS-A 0499	2.9e-06	4.9e-04	2.1e-04	
New Jersey OCS-A 0498	2.8e-06	5.3e-04	2.1e-04	
Delaware OCS-A 0482	3.3e-06	7.4e-04	1.3e-04	
Maryland OCS-A 0489	2.1e-06	2.5e-04	5.7e-05	
Maryland OCS-A 0490	2.2e-06	1.9e-04	5.0e-05	
Virginia OCS-A 0483	3.9e-06	9.1e-05	1.0e-04	
Virginia OCS-A 0497	3.2e-06	4.4e-05	8.4e-06	
North Carolina OCS-A 0508	3.4e-06	3.2e-05	8.8e-05	
North Carolina WEA - Wilmington West	9.7e-04	6.9e-04	8.4e-04	
North Carolina WEA - Wilmington East	2.1e-04	6.4e-04	6.0e-04	
South Carolina Call Area - Grand Strand	1.0e-02	1.1e-02	9.1e-03	
South Carolina Call Area - Cape Romain	9.1e-03	4.6e-03	6.3e-03	
South Carolina Call Area - Winyah	4.4e-04	1.2e-05	6.4e-05	
South Carolina Call Area - Charleston	3.7e-05	4.5e-05	7.5e-05	
All	2.1e-02	1.9e-02	1.8e-02	

Predicted relative abundance of RTLO as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	9.3e-04		3.3e-04	2.9e-04
Rhode Island / Massachusetts OCS-A 0486	6.6e-04		4.1e-04	4.6e-04
Rhode Island / Massachusetts OCS-A 0487	3.1e-04		2.2e-04	1.9e-04
Massachusetts OCS-A 0500	8.1e-04		5.3e-04	4.9e-04
Massachusetts OCS-A 0501	8.6e-04		4.5e-04	4.4e-04
Massachusetts OCS-A 0502	9.3e-04		5.3e-04	6.0e-04
Massachusetts OCS-A 0503	4.3e-04		2.6e-04	3.6e-04
New York Proposed Commercial Lease - Unsolicited	2.0e-04		1.4e-04	1.6e-04
New York OCS-A 0512	6.0e-04		3.2e-04	6.7e-04
New Jersey OCS-A 0499	5.1e-03		8.3e-04	2.0e-03
New Jersey OCS-A 0498	3.0e-03		6.9e-04	1.9e-03
Delaware OCS-A 0482	1.3e-03		3.8e-04	1.2e-03
Maryland OCS-A 0489	4.4e-04		1.4e-04	5.1e-04
Maryland OCS-A 0490	4.9e-04		1.6e-04	5.2e-04
Virginia OCS-A 0483	9.2e-04		2.8e-04	1.0e-03
Virginia OCS-A 0497	4.6e-05		8.4e-06	4.8e-05
North Carolina OCS-A 0508	4.4e-04		2.7e-04	8.9e-04
North Carolina WEA - Wilmington West	2.1e-03		1.8e-04	6.5e-04
North Carolina WEA - Wilmington East	6.4e-04		3.4e-04	8.7e-04
South Carolina Call Area - Grand Strand	1.9e-02		2.1e-03	7.2e-03
South Carolina Call Area - Cape Romain	2.3e-03		6.1e-04	1.6e-03
South Carolina Call Area - Winyah	7.0e-05		8.2e-05	8.1e-05
South Carolina Call Area - Charleston	9.9e-05		9.1e-05	1.6e-04
All	4.1e-02		9.4e-03	2.2e-02

Predicted relative abundance of SOSH as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	3.5e-05	1.2e-05	4.6e-05	
Rhode Island / Massachusetts OCS-A 0486	2.0e-04	4.4e-04	1.4e-04	
Rhode Island / Massachusetts OCS-A 0487	1.9e-04	5.1e-04	9.6e-05	
Massachusetts OCS-A 0500	5.9e-04	8.3e-04	2.7e-04	
Massachusetts OCS-A 0501	4.6e-04	7.0e-04	2.4e-04	
Massachusetts OCS-A 0502	8.9e-04	1.0e-03	3.5e-04	
Massachusetts OCS-A 0503	5.2e-04	6.6e-04	2.0e-04	
New York Proposed Commercial Lease - Unsolicited	6.8e-05	7.1e-05	5.5e-05	
New York OCS-A 0512	9.9e-05	1.1e-04	1.1e-04	
New Jersey OCS-A 0499	1.5e-04	5.9e-05	2.3e-04	
New Jersey OCS-A 0498	1.4e-04	4.7e-05	2.1e-04	
Delaware OCS-A 0482	7.9e-05	2.5e-05	1.2e-04	
Maryland OCS-A 0489	2.3e-05	8.2e-06	4.1e-05	
Maryland OCS-A 0490	3.8e-05	1.2e-05	5.8e-05	
Virginia OCS-A 0483	1.1e-04	2.5e-05	1.4e-04	
Virginia OCS-A 0497	2.2e-06	6.2e-07	3.3e-06	
North Carolina OCS-A 0508	2.0e-04	2.7e-05	1.5e-04	
North Carolina WEA - Wilmington West	2.9e-05	1.1e-05	5.8e-05	
North Carolina WEA - Wilmington East	8.6e-05	2.8e-05	1.5e-04	
South Carolina Call Area - Grand Strand	3.6e-04	1.3e-04	7.1e-04	
South Carolina Call Area - Cape Romain	8.1e-05	3.4e-05	1.8e-04	
South Carolina Call Area - Winyah	2.4e-05	7.1e-06	4.4e-05	
South Carolina Call Area - Charleston	2.3e-05	7.3e-06	4.0e-05	
All	4.4e-03	4.8e-03	3.6e-03	

Predicted relative abundance of SOTE as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	5.9e-16	1.9e-06		
Rhode Island / Massachusetts OCS-A 0486	2.0e-15	6.3e-06		
Rhode Island / Massachusetts OCS-A 0487	1.3e-15	4.4e-06		
Massachusetts OCS-A 0500	3.8e-15	1.2e-05		
Massachusetts OCS-A 0501	3.3e-15	1.1e-05		
Massachusetts OCS-A 0502	5.0e-15	1.6e-05		
Massachusetts OCS-A 0503	2.8e-15	9.0e-06		
New York Proposed Commercial Lease - Unsolicited	8.2e-16	2.8e-06		
New York OCS-A 0512	1.6e-15	5.4e-06		
New Jersey OCS-A 0499	3.6e-15	1.3e-05		
New Jersey OCS-A 0498	3.2e-15	1.2e-05		
Delaware OCS-A 0482	1.9e-15	7.2e-06		
Maryland OCS-A 0489	6.5e-16	2.4e-06		
Maryland OCS-A 0490	9.4e-16	3.5e-06		
Virginia OCS-A 0483	2.3e-15	9.2e-06		
Virginia OCS-A 0497	5.6e-17	2.4e-07		
North Carolina OCS-A 0508	2.5e-15	1.0e-05		
North Carolina WEA - Wilmington West	1.1e-15	9.7e-06		
North Carolina WEA - Wilmington East	3.1e-15	1.7e-03		
South Carolina Call Area - Grand Strand	1.4e-14	2.1e-04		
South Carolina Call Area - Cape Romain	3.3e-15	8.3e-05		
South Carolina Call Area - Winyah	7.4e-16	5.1e-05		
South Carolina Call Area - Charleston	7.4e-16	2.2e-05		
All	5.9e-14	2.2e-03		

Predicted relative abundance of SPSK as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478		3.6e-05	1.2e-05	
Rhode Island / Massachusetts OCS-A 0486		1.2e-04	3.6e-05	
Rhode Island / Massachusetts OCS-A 0487		8.8e-05	2.6e-05	
Massachusetts OCS-A 0500		2.6e-04	7.7e-05	
Massachusetts OCS-A 0501		2.5e-04	7.1e-05	
Massachusetts OCS-A 0502		4.0e-04	1.1e-04	
Massachusetts OCS-A 0503		2.7e-04	7.2e-05	
New York Proposed Commercial Lease - Unsolicited		4.4e-05	1.2e-05	
New York OCS-A 0512		7.8e-05	2.2e-05	
New Jersey OCS-A 0499		1.8e-04	4.9e-05	
New Jersey OCS-A 0498		1.5e-04	4.3e-05	
Delaware OCS-A 0482		9.1e-05	2.6e-05	
Maryland OCS-A 0489		3.1e-05	8.8e-06	
Maryland OCS-A 0490		4.5e-05	1.3e-05	
Virginia OCS-A 0483		1.1e-04	3.3e-05	
Virginia OCS-A 0497		2.6e-06	7.8e-07	
North Carolina OCS-A 0508		1.3e-04	5.9e-05	
North Carolina WEA - Wilmington West		4.3e-05	1.2e-05	
North Carolina WEA - Wilmington East		1.2e-04	3.3e-05	
South Carolina Call Area - Grand Strand		5.1e-04	1.4e-04	
South Carolina Call Area - Cape Romain		1.2e-04	3.3e-05	
South Carolina Call Area - Winyah		3.1e-05	8.8e-06	
South Carolina Call Area - Charleston		2.9e-05	7.9e-06	
All		3.1e-03	9.1e-04	

Predicted relative abundance of SUSC as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	6.4e-03		7.6e-03	3.7e-03
Rhode Island / Massachusetts OCS-A 0486	8.1e-05		2.2e-04	2.5e-04
Rhode Island / Massachusetts OCS-A 0487	4.2e-05		1.3e-04	1.6e-04
Massachusetts OCS-A 0500	1.3e-04		3.5e-04	4.7e-04
Massachusetts OCS-A 0501	1.7e-04		3.1e-04	4.3e-04
Massachusetts OCS-A 0502	4.0e-04		4.4e-04	6.7e-04
Massachusetts OCS-A 0503	2.7e-04		2.4e-04	4.0e-04
New York Proposed Commercial Lease - Unsolicited	2.6e-05		8.8e-05	1.1e-04
New York OCS-A 0512	1.5e-04		2.8e-04	2.3e-04
New Jersey OCS-A 0499	1.0e-02		6.2e-04	5.3e-04
New Jersey OCS-A 0498	3.3e-04		6.0e-04	4.9e-04
Delaware OCS-A 0482	1.8e-04		3.2e-04	3.3e-04
Maryland OCS-A 0489	7.7e-05		1.2e-04	1.2e-04
Maryland OCS-A 0490	5.4e-05		1.1e-04	1.4e-04
Virginia OCS-A 0483	7.2e-05		1.9e-04	3.2e-04
Virginia OCS-A 0497	4.5e-05		5.8e-06	1.9e-05
North Carolina OCS-A 0508	7.0e-05		1.9e-04	3.1e-04
North Carolina WEA - Wilmington West	9.6e-05		1.0e-04	1.4e-04
North Carolina WEA - Wilmington East	9.0e-05		2.0e-04	3.1e-04
South Carolina Call Area - Grand Strand	1.6e-03		1.2e-03	1.6e-03
South Carolina Call Area - Cape Romain	7.6e-04		3.3e-04	4.5e-04
South Carolina Call Area - Winyah	1.7e-05		5.2e-05	7.6e-05
South Carolina Call Area - Charleston	1.8e-05		5.5e-05	7.8e-05
All	2.1e-02		1.4e-02	1.1e-02

Predicted relative abundance of TBMU as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	1.6e-05			1.3e-05
Rhode Island / Massachusetts OCS-A 0486	4.4e-05			4.4e-05
Rhode Island / Massachusetts OCS-A 0487	3.1e-05			3.1e-05
Massachusetts OCS-A 0500	8.9e-05			1.0e-04
Massachusetts OCS-A 0501	8.3e-05			9.4e-05
Massachusetts OCS-A 0502	1.3e-04			1.6e-04
Massachusetts OCS-A 0503	1.0e-04			9.1e-05
New York Proposed Commercial Lease - Unsolicited	1.4e-05			1.6e-05
New York OCS-A 0512	2.6e-05			3.0e-05
New Jersey OCS-A 0499	5.4e-05			6.4e-05
New Jersey OCS-A 0498	4.5e-05			5.9e-05
Delaware OCS-A 0482	2.6e-05			3.5e-05
Maryland OCS-A 0489	8.3e-06			1.3e-05
Maryland OCS-A 0490	1.2e-05			1.8e-05
Virginia OCS-A 0483	2.7e-05			4.8e-05
Virginia OCS-A 0497	6.6e-07			1.2e-06
North Carolina OCS-A 0508	3.0e-05			6.4e-05
North Carolina WEA - Wilmington West	1.0e-05			2.2e-05
North Carolina WEA - Wilmington East	2.6e-05			7.0e-05
South Carolina Call Area - Grand Strand	1.2e-04			2.6e-04
South Carolina Call Area - Cape Romain	2.9e-05			5.9e-05
South Carolina Call Area - Winyah	1.2e-05			1.8e-05
South Carolina Call Area - Charleston	7.2e-06			1.7e-05
All	9.4e-04			1.3e-03

Predicted relative abundance of WISP as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	2.2e-05	2.5e-05	1.8e-05	
Rhode Island / Massachusetts OCS-A 0486	6.7e-05	5.5e-04	1.0e-04	
Rhode Island / Massachusetts OCS-A 0487	5.2e-05	9.1e-04	1.1e-04	
Massachusetts OCS-A 0500	1.6e-04	1.7e-03	2.5e-04	
Massachusetts OCS-A 0501	1.4e-04	1.3e-03	1.8e-04	
Massachusetts OCS-A 0502	3.7e-04	1.9e-03	3.2e-04	
Massachusetts OCS-A 0503	2.6e-04	9.3e-04	2.0e-04	
New York Proposed Commercial Lease - Unsolicited	2.8e-05	2.9e-04	5.4e-05	
New York OCS-A 0512	4.9e-05	3.4e-04	9.5e-05	
New Jersey OCS-A 0499	1.2e-04	5.2e-04	1.6e-04	
New Jersey OCS-A 0498	1.0e-04	4.8e-04	1.1e-04	
Delaware OCS-A 0482	6.3e-05	2.3e-04	5.0e-05	
Maryland OCS-A 0489	2.2e-05	5.4e-05	1.6e-05	
Maryland OCS-A 0490	3.2e-05	9.2e-05	2.6e-05	
Virginia OCS-A 0483	1.0e-04	2.1e-04	6.9e-05	
Virginia OCS-A 0497	2.1e-06	4.1e-06	1.3e-06	
North Carolina OCS-A 0508	1.7e-04	3.2e-04	1.4e-04	
North Carolina WEA - Wilmington West	3.2e-05	2.2e-05	2.3e-05	
North Carolina WEA - Wilmington East	8.8e-05	5.7e-05	6.0e-05	
South Carolina Call Area - Grand Strand	3.8e-04	2.5e-04	2.8e-04	
South Carolina Call Area - Cape Romain	9.4e-05	5.6e-05	6.8e-05	
South Carolina Call Area - Winyah	3.0e-05	1.4e-05	2.4e-05	
South Carolina Call Area - Charleston	2.4e-05	1.4e-05	1.6e-05	
All	2.4e-03	1.0e-02	2.4e-03	

Predicted relative abundance of WWSC as a proportion of total relative abundance in the study area by BOEM area (Fig. D1) and season. Note that these results are relative to the study area. They do not account for birds outside of the study area. If the entire range of a species were to be considered then the proportional relative abundance in BOEM areas could decrease.

BOEM area	Spring	Summer	Fall	Winter
Massachusetts OCS-A 0478	4.9e-04		9.9e-03	4.0e-03
Rhode Island / Massachusetts OCS-A 0486	2.8e-04		2.0e-04	8.3e-04
Rhode Island / Massachusetts OCS-A 0487	1.8e-04		1.0e-04	2.7e-04
Massachusetts OCS-A 0500	5.2e-04		2.6e-04	9.8e-04
Massachusetts OCS-A 0501	4.7e-04		2.4e-04	1.3e-03
Massachusetts OCS-A 0502	7.9e-04		3.3e-04	1.8e-03
Massachusetts OCS-A 0503	3.1e-03		1.8e-04	4.9e-03
New York Proposed Commercial Lease - Unsolicited	1.1e-04		5.9e-05	1.5e-04
New York OCS-A 0512	2.2e-04		1.3e-04	4.1e-04
New Jersey OCS-A 0499	5.4e-04		2.7e-04	8.5e-04
New Jersey OCS-A 0498	4.2e-04		2.2e-04	7.0e-04
Delaware OCS-A 0482	2.5e-04		1.3e-04	3.4e-04
Maryland OCS-A 0489	8.8e-05		4.8e-05	1.1e-04
Maryland OCS-A 0490	1.2e-04		5.9e-05	1.3e-04
Virginia OCS-A 0483	2.8e-04		1.3e-04	2.5e-04
Virginia OCS-A 0497	7.7e-06		3.1e-06	6.5e-06
North Carolina OCS-A 0508	3.0e-04		1.4e-04	2.7e-04
North Carolina WEA - Wilmington West	1.4e-04		6.1e-05	9.9e-05
North Carolina WEA - Wilmington East	3.3e-04		1.5e-04	2.6e-04
South Carolina Call Area - Grand Strand	1.7e-03		7.5e-04	1.2e-03
South Carolina Call Area - Cape Romain	4.3e-04		1.8e-04	3.2e-04
South Carolina Call Area - Winyah	8.5e-05		3.8e-05	7.3e-05
South Carolina Call Area - Charleston	8.7e-05		3.8e-05	6.9e-05
All	1.1e-02		1.4e-02	1.9e-02